



CLINICAL RESEARCH:

Prevalence of Lesions and Oral Mucosal Normal Variations in an Elderly Population in Costa Rica

Prevalencia de lesiones y variaciones de lo normal en la mucosa oral en una población de adultos mayores de Costa Rica

José Manuel Fernández-Chaves DDS, MSc¹ <https://orcid.org/0000-0001-6478-5407>

Yadira V. Boza-Oreamuno DDS, MSc¹ <https://orcid.org/0000-0002-0367-8664>

Carlos Masís-Monestel DDS² <https://orcid.org/0000-0003-0303-6343>

Norma Lau-Sánchez BSN, MSc³ <https://orcid.org/0000-0002-8426-3101>

1. Department of Diagnostic and Surgical Sciences, Faculty of Dentistry, University of Costa Rica, San José, Costa Rica.
2. Department of Social Dentistry, Faculty of Dentistry, University of Costa Rica, San José, Costa Rica.
3. Instituto de Investigaciones en Salud, University of Costa Rica, San José, Costa Rica.

Correspondence to: Dr. José Manuel Fernández Chaves- jose.fernandezchaves@ucr.ac.cr

Received: 8-VII-2023

Accepted: 1-X-2023

ABSTRACT: To establish the prevalence of oral mucosal lesions and normal variations in a population of older adults. This observational study was conducted from 2015 to 2016 and involved 156 participants from Los Guido (San José, Costa Rica). The participants' sociodemographic information, risk factors, comorbidity, and clinical examination were evaluated and analyzed descriptively and qualitatively. Findings in oral mucosa were found to preferentially affect women (n=111; 71%) and individuals between the ages of 60 and 69 years (n=65; 42%), having smoking as a risk factor (n=67; 43%), and suffering from arterial hypertension (n=101; 65%). Intraoral findings were mainly presented together with variations from normal and pathology (n=71; 46%), predominantly one variation from normal (n=67; 33%) and one pathology (n=51; 43%). The most prevalent normal variations were the saburral tongue, lingual varicose veins, and cryptic tonsils. Infectious diseases (n=33; 21%), potentially malignant disorders (n=30; 19%), and physical and chemical injuries (n=18; 12%) were the most prevalent groups. The four most predominant lesions were prosthetic stomatitis, actinic cheilitis, frictional keratosis, and atrophic glossitis. The differences in pathological groups were not statistically significant (p=0.392) when comparing the frequency ratio by sex. Variations of the normal anatomy are the predominant ones, the saburral tongue being the most prevalent individually; among the pathologies, the most remarkable group was the infectious diseases, and prosthetic stomatitis was the most frequent. The information



about this subject in Costa Rica is limited. So it is important to conduct more studies that contribute to the oral health of the elderly population in this area.

KEYWORDS: Oral cancer screening; Oral mucosa; Aging; Elderly; Oral lesions; Normal variations.

RESUMEN: Establecer la prevalencia de lesiones de la mucosa oral y variaciones de lo normal en una población de adultos mayores. Estudio observacional en 156 participantes de Los Guido (San José, Costa Rica) del 2015 a 2016. Se evaluaron y analizaron de forma descriptiva y cualitativa información sociodemográfica, factores de riesgo, comorbilidad y exploración clínica. Los hallazgos en mucosa oral afectaron preferentemente a las mujeres (n=111; 71%), el rango de edad entre 60 y 69 años (n=65; 42%) y con el tabaco como factor de riesgo (n=67; 43%) e hipertensión arterial (n=101; 65%). Los hallazgos intraorales se presentaron preferentemente de forma conjunta variaciones de lo normal y patologías (n=71; 46%), predominantemente una variación de lo normal (n=67; 33%) y una patología (n=51; 43%). Las variaciones de lo normal más prevalentes fueron lengua saburral, várices linguales y amígdalas criptodónticas. Las enfermedades infecciosas (n=33; 21%), los desórdenes potencialmente malignos (n=30; 19 %) e injurias físicas y químicas (n=18; 12%) fueron los grupos más prevalentes. Las cuatro lesiones más predominantes fueron estomatitis protésica, queilitis actínica, queratosis friccional y glositis atrófica. No hubo diferencia estadísticamente significativa ($p=0.392$) al comparar la proporción de frecuencia por sexo. Las variaciones de la anatomía normal predominaron; siendo en forma individual la lengua saburral la más frecuente; de las patologías destacaron las enfermedades infecciosas principalmente la estomatitis protésica. Existe poca información en esta área en Costa Rica, es importante generar más estudios que permitan contribuir con la salud oral de la población adulta mayor.

PALABRAS CLAVE: Tamizaje de cáncer bucal; Mucosa oral; Envejecimiento; Lesiones orales; Variaciones de lo normal.

INTRODUCTION

Human aging is a complex multifactorial process that leads to gradually deteriorating bodily functions, productivity, and psychological changes, as well as multiple diseases that inevitably end in death. However, certain lifestyle and behavioral modifications can help slow this decline and keep the body in the best possible state for its chronological age, commonly referred to as “healthy aging” (1).

By 2050, one in six people in the world (16% of the population) will be over 65 years old, compared with one in 11 people in 2019 (9%). The proportion of the population aged 65 and over is projected to double between 2019 and 2050; in 2018, for the first time in history, people aged 65 and over outnumbered children under five years old globally (2).

In Costa Rica, the population over 65 years of age constituted 6% of the total population in

2008 and 8.9% in 2020, and according to the population projections of the National Institute of Statistics and Censuses (INEC), it is estimated to reach 13% by 2030 and more than 20% in 2050 (3).

The oral cavity does not escape changes brought on by aging. Many of these changes are manifestations of pathologies such as immunological diseases, hematological disorders, or other diseases such as diabetes mellitus or nutritional deficiencies (4). Other changes may be associated with local factors such as the use of dentures, as frequently found in other studies (4,5). Most oral mucosal lesions are benign, the most common being fibrous hyperplasia (6,7); among malignant tumors, squamous cell carcinoma (SCC) is the most common pathology (8).

In recent years, research has been conducted worldwide on the epidemiological aspects of oral mucosal lesions (9,10); however, few studies have been carried out in Costa Rica (7,11-14).

A complete and systematic clinical oral assessment is essential for detecting any lesion or variation. This assessment must be performed in the same order in each of the evaluated patients to ensure that no findings are omitted, including palpation of nodes and review of each of the structures of the oral cavity, starting with the lips (15). All observations must be properly documented in the dental record per the legal requirements of each country (16,17).

In this context, the objective of the current study was to determine the prevalence of mucosal normal variations and lesions in a population of older adults from Los Guido de Desamparados (San José, Costa Rica), using the socio-demographic data in the period from 2015 to 2016.

MATERIALS AND METHODS

PARTICIPANTS

The research was conducted for the period from 2015 to 2016. Elderly people were selected by systematic random sampling with the participants' consent. A sample size was selected according to the convenience of the researchers. The inclusion criteria were patients aged 60 years or older from the Los Guido de Desamparados area of Costa Rica who had agreed to participate in the research. The exclusion criteria were patients who did not belong to the indicated area, refused to be examined, had a situation that prevented them from revealing their clinical history, or could not be examined.

ETHICAL CONSIDERATIONS

The Scientific Ethics Committee of the University of Costa Rica (UCR) approved this descriptive observational study under project code B2335. The patients gave their free and informed consent in writing to participate in the study. The confidentiality and anonymization of the collected data were ensured.

DATA COLLECTION

According to the INEC-prepared census maps, a random area was selected within the Los Guido district to ensure that all communities had an equal chance of inclusion in the study. The study team registered and selected households in each community that met the inclusion criteria until the sample was complete. Those who agreed to participate were invited to the same location for an oral cancer screening, educational talk, and intraoral examination. The team also set up a mobile

clinic for history-taking, intraoral examination, and ganglion palpation (See Figure 1 for details).

The variables studied were age (completed years); sex of the patient (male, female); history of

cancer (personal and family); risk factors (tobacco, alcohol, and sexual practices); comorbidity (systemic diseases); and edentulism. For oral mucosal findings, variations from normal and pathologies according to Neville's classification were considered (18).




Figure 1. Diagram to carry out oral cancer screening. Source: self-made.

STATISTICAL ANALYSIS

The data were entered in an Excel spreadsheet (Microsoft, Inc., Redmond, WA, USA). Inconsistencies were entered, corrected, and subsequently analyzed using descriptive statistics and the Chi-square test with the Stata 14 program (Statistical software package, Stata Corp). The significance level of the statistical difference was $p\text{-value} < 0.05$, with a confidence interval of 95%.

RESULTS

A total of 156 patients were examined. Most of the participants were female (71%), 42% of them were between 60 and 69 years of age, 43% had smoking as a risk factor, 65% had reported a history of arterial hypertension, and 44% had diabetes mellitus as comorbidities. Additionally, 15% of the participants had palpable lymph nodes (Table 1).

Table 1. Distribution of sociodemographic variables (N = 156).

Variable	N	%
Sex		
Male	45	29%
Female	111	71%
Age		
60–69 years	65	42%
70–79 years	59	38%
80 years and over	32	21%
History of cancer		
Personal history of cancer	5	3%
Family history of cancer	12	8%
Risk factors		
Alcohol	7	4%
Tobacco	67	43%
Risky sexual practices	23	15%
Comorbidity		
Arterial hypertension	101	65%
Diabetes mellitus	69	44%
Asthma	26	17%
Hypothyroidism	24	15%
Arthritis	3	2%
Gastritis	15	10%
Anemia	21	13%
Other	49	31%
Palpation of nodes		
Yes	24	15%
No	129	83%
Not reported	3	2%

Source: self-made.

The intraoral examination findings most frequently found were both normal variations and pathology (46%): about 43% of the participants had only one variation from normal, while 33% exhibited one pathology (Table 2).

The most prevalent normal variations were saburral tongue (54%), lingual varicose veins (26%), and cryptic tonsils (15%). Pathologies were classified into 10 groups, of which infectious diseases (21%), potentially malignant disorders (PMD) (19%), and physical and chemical injuries

(12%) were the most prevalent. Individually, most findings comprised the diagnosis of prosthetic stomatitis (24 cases) comprised. The second- and third-most prevalent diagnoses in this study were actinic cheilitis (23 cases) and frictional keratosis (10 cases) and atrophic glossitis (10 cases), respectively (Table 3).

When comparing the frequency ratio by sex in these groups of lesions, the difference in the distribution for either group was not statistically significant ($p=0.392$; Table 4).

Table 2. Frequency of findings in the intraoral clinical examination.

	N	%
Type of finding		
None	12	8%
Normal variations	55	35%
Pathological	18	12%
Normal variations and pathological	71	46%
Number of variations		
1	67	43%
2	48	31%
3	11	7%
Number of pathologies		
1	51	33%
2	31	20%
3	7	4%
Total	156	

Source: self-made.

Table 3. Frequency of variations from normal and oral lesions.

Variable	N	%
Normal variations		
Torus	3	2%
Fibroepithelial polyp	0	0%
Lingual varicosities	40	26%
Hypertrophic lymphoid tissue	0	0%
Cryptic (hypertrophic) tonsils	23	15%
Fordyce granules	14	9%
Leukoedema	4	3%
Macroglossia	0	0%
Linea alba	0	0%
Physiological pigmentation	13	8%
Fissured tongue	13	8%
Saburral (coated) tongue	84	54%
Ankyloglossia	2	1%
Total	196	100%
Group of pathologies		
Non-neoplastic proliferative lesions		
Fibrous hyperplasia (traumatic fibroma)	5	3%
Inflammatory fibrous hyperplasia (epulis)	6	4%
Potentially malignant disorders		
Actinic cheilitis	23	15%
Leukoplakia	6	4%
Erythroplakia	3	2%
Salivary gland lesions		
Mucocele	1	1%
Physical and chemical injuries		
Traumatic ulcer	5	3%
Frictional keratosis	10	6%
Petechiae and ecchymosis	3	2%
Pigmented lesions		
Melanotic macule	7	4%
Nevus	1	1%
Smoker melanosis	1	1%
Amalgam tattoo	1	1%
Infectious diseases		
Erythematous candidiasis	1	1%
Denture stomatitis	24	15%
Median rhomboid glossitis	5	3%
Angular cheilitis	4	3%
Pseudomembranous candidiasis	2	1%

Variable	N	%
Vascular lesions	1	1%
Hemangioma	1	1%
Benign mesenchymal lesions	1	1%
Lipoma	1	1%
Periodontal disease	5	3%
Periodontal disease	5	3%
Unrated	19	12%
Tonsillolith	1	1%
Nicotinic stomatitis	0	0%
Atrophic glossitis	10	6%
Aphthae	4	3%
Xerostomia (dry mouth)	2	1%
Total	134	100%

Source: self-made.

Table 4. Frequency by sex in the groups according to clinical diagnosis of oral pathologies.

Group	N	% total	Sex		Half of age (standard derivation)
			Male (%)	Female (%)	
Non-neoplastic proliferative lesions	11	7%	2%	9%	71
Potentially malignant disorders	30	19%	11%	23%	73
Salivary gland lesions	1	1%	0%	1%	-
Physical and chemical injuries	18	12%	7%	14%	77
Pigmented lesions	10	6%	9%	5%	73
Infectious diseases	33	21%	13%	24%	72
Vascular lesions	1	1%	2%	0%	-
Benign mesenchymal lesions	1	1%	2%	0%	-
Periodontal disease	5	3%	7%	2%	73
Unrated	19	12%	11%	13%	73

**Chi-square test $p = 0.392$.

Source: self-made.

DISCUSSION

The aging of the population is considered one of the greatest challenges in the world; by 2030, it is estimated that all baby boomers will be over 65 years of age. Hence, knowing the etiological factors, pathogenesis, and coadjutant factors that determine the specificity of oral conditions in this age group is important (19).

The present study found that females exhibited oral mucosal findings more frequently than males did. Other studies in the literature have also reported this trend (20-23). This may be explained by the fact that multiple studies have found that women use health services more frequently than men do, which increases the possibility of detection, as observed in countries such as Brazil (24-26), Chile (27), Mexico (28), Italy (29), Poland (4), and Spain (30).

The most prevalent normal variations were saburrall tongue and lingual varicose veins. Additionally, our study population had several causal factors reported in the literature. Saburrall tongue has been associated with poor oral hygiene, dehydration, tobacco use, and alcohol consumption (31); the main systemic disease linked to Saburrall tongue is arterial hypertension followed by diabetes mellitus, gastrointestinal disorders, and arthritis (32). Aging, smoking, and systemic diseases such as arterial hypertension, cardiovascular or cardiopulmonary diseases, and diabetes mellitus, have been associated with sublingual varicose veins (4).

Lesions associated with infectious diseases were the most prevalent group, affecting women more frequently; prosthetic stomatitis was the most common diagnosis in the entire sample, followed by different presentations of oral candidiasis (34) in several other countries according to a systema-

tic review (34). Notably, the aging process produces gradual, irreversible, and cumulative changes in the oral cavity, increasing its vulnerability to traumatic and infectious agents (35).

These lesions are influenced by changes in salivary function and oral microbiota, the quantity and composition of saliva change due to aging, systemic diseases, and medications used. Bacteria in older adults have been proven to change from being predominantly aerobic gram-positive to gram-negative over time. This transformation, along with changes in the mucosa, affects the immunological activity in the oral cavity, making it susceptible to infections by opportunistic microorganisms (19).

The second-most prevalent group in the lesions was PMD, clinically represented mostly by actinic cheilitis. Other studies have also reported these findings (21, 36), but this result is highly relevant as it has been concluded that the possibility of presenting with PMD and malignant lesions increases with age (33) and that retrospective studies have found SCC to be the most common lesion along with a high frequency of potentially malignant oral disorders (21, 37, 38).

Actinic cheilitis occurs between 4.6 to 43.2% of the worldwide population. Populations in tropical regions with excessive exposure to ultraviolet rays are at higher risk (39). A multicenter Spanish study showed that among the risk factors in the multivariate analysis were individuals being older than 60 years, having worked under sun exposure for more than 25 years, and having a history of skin cancer (non-melanoma) (40). The population studied in Los Guido exhibited these factors in the majority of cases.

The third-most prevalent diagnosis was frictional keratosis and atrophic glossitis, both of

which are reported as frequent findings in adult denture wearers in studies conducted in Poland (4), Chile (41), and Brazil (42). Atrophic glossitis is most likely a pernicious anemia manifestation (43). However, no hematological evidence supports this.

These findings are related to normal physiological processes during aging, in which the oral mucosa presents decreased elastic fibers and thinned and disorganized collagen fibers in the connective tissue, as well as decreased microvasculature. Due to these processes, the mucosa is less resilient and has reduced healing capacity and, with the previously mentioned microbiota changes, reduced immune response (19).

In this project, an educational talk was delivered to all participants in a community hall, and during the screening and assessment, the concepts of prevention and detection of oral cancer were explained to them individually. Previous studies have shown that patients receive more information from the media than from dental and medical professionals (44). Educational campaigns generate an increase in not only knowledge but also public awareness and thus early diagnosis (45).

In Costa Rica, a study conducted in 2004 showed that only 10% of the patients interviewed reported having undergone a head and neck examination by dental professionals, and an even lower percentage (9.7%) mentioned other oral tissues in the mouth being examined besides teeth and gums (46).

The presence of oral lesions is not always obvious and is not necessarily accompanied by symptoms, but many can be detected early through routine screening (47), including inspection and palpation of the neck and oral cavity

using a bimanual technique (48). This technique, in addition to being economical and simple, does not cause discomfort to patients (49). The World Health Organization recommends screening as it can reduce mortality in oral cancer patients (50).

Oral cavity examination should be routinely performed in older adults every year as a preventive measure that can reduce morbidity and mortality in this population (35). Screening also detects conditions that, although not malignant, can damage a patient's oral health, as in our case with prosthetic stomatitis, atrophic glossitis, and frictional keratosis. In short, knowing the normal variations of the oral mucosa avoids unnecessary procedures.

Developing specific oral health policies is necessary to address the needs of older adults with programs sustainable over time to allow this population to maintain adequate nutrition and detect lesions early (51).

The main limitation in this project was scheduling the visits as the hours of service had to be before noon because the community is at social risk and its location endangers the staff. Second, the lack of infrastructure at the site demanded the use of portable equipment in less-than-ideal conditions for data collection. Moreover, the health condition of many elderly people prevented them from moving, so the professional team had to visit their homes, which generated greater insecurity. Finally, the health condition of the participating population caused absenteeism at evaluation appointments, added to the multiple activities and functions of each researcher, and made it difficult to coordinate the dates together with the security conditions, which on some occasions forced the cancellation of the tours due to criminal acts or conflicts the day before.

The results of this study allow not only the Faculty of Dentistry of the UCR but also other public and private entities to design and program oral health activities aimed at the elderly population to reduce the risk factor for oral lesions, as well as to investigate, design, and evaluate the quality of life related to oral health for this population group that will inevitably increase in Costa Rica in the coming years.

CONCLUSIONS

According to oral screenings performed on a population of older adults in Los Guido de Desamparados in Costa Rica from 2015 to 2016, this study found that the variations of normal anatomy were predominant, the saburral tongue being the most prevalent individually; among the pathologies, the most prevalent groups were the infectious diseases, while prosthetic stomatitis was the most frequent.

There is little information about this subject in Costa Rica; therefore, it is important to conduct more studies to contribute to the oral health of the elderly population.

CONFLICTS OF INTEREST

The authors declare no conflict of interest.

FUNDING

University of Costa Rica Vice Rector's Office for Research ordinary funds Project B2335 given to MSc. Norma Lau Sánchez.

AUTHOR CONTRIBUTION STATEMENT

Conceptualization: Y.V.B.O. and J.M.F.CH.
Methodology: Y.V.B.O. and J.M.F.CH.
Validation: Y.V.B.O. and J.M.F.CH.
Formal analysis: Y.V.B.O. and J.M.F.CH.
Investigation: Y.V.B.O. and J.M.F.CH.
Resources: Y.V.B.O., J.M.F.CH., N.L.S. and C.M.M.
Data curation: Y.V.B.O. and J.M.F.CH.
Writing-original draft preparation: Y.V.B.O. and J.M.F.CH.
Writing-review and editing: Y.V.B.O. and J.M.F.CH.
Visualization: Y.V.B.O. and J.M.F.CH.
Project administration: N.L.S.
Funding acquisition: N.L.S.

All authors have read and agreed with the published version of the manuscript.

ACKNOWLEDGMENTS

The authors acknowledge M.Sc. Romain Fantin for collaborating in the statistical analysis of the data and Kattia Campos Chaves from the Graphic Design Area of the Multimedia Unit of the Faculty of Dentistry of the University of Costa Rica for editing Figure 1.

Special thanks to Dr. Madeline Howard Mora for her invaluable collaboration with this project.

REFERENCES

1. Zhavoronkov A., Li R., Ma C., Mamoshina P. Deep biomarkers of aging and longevity: from research to applications. *Aging (Albany NY)*. 2019 Nov 11; 11 (22): 10771.

2. United Nations. World Population Prospects: The 2022 Revision. Methodology of the United Nations Population Estimates and Projections. *Econ Soc Aff.* 2022; 6 (July 2022): 54.
3. Demográficamente, población costarricense se encuentra en proceso de envejecimiento | Ministerio de Planificación Nacional y Política Económica [Internet]. [cited 2022 Oct 16]. Available from: <https://www.mideplan.go.cr/demograficamente-poblacion-costarricense-se-encuentra-en-proceso-de-envejecimiento>
4. Radwan-Oczko M., Bandosz K., Rojek Z., Owczarek-Drabińska J.E. Clinical Study of Oral Mucosal Lesions in the Elderly-Prevalence and Distribution. *Int J Environ Res Public Health.* 2022 Mar 1; 19 (5): 1-10.
5. Marcia Hortensia Corona Carpio D., Dra Ruth Ramón Jimenez I., Wiljer Urgellés Matos I. Lesiones de la mucosa bucal en adultos mayores con prótesis dentales totales. *Medisan.* 2017; 21 (7): 813-8.
6. Maturana-Ramírez A., Adorno-Farías D., Reyes-Rojas M., Farías-Vergara M., Aitken-Saavedra J. A retrospective analysis of reactive hyperplastic lesions of the oral cavity: study of 1149 cases diagnosed between 2000 and 2011, Chile. *Acta Odontol Latinoam.* 2015; 28 (2): 103-7.
7. Boza Oreamuno Y., López Soto A. Análisis retrospectivo de las lesiones de la mucosa oral entre 2008-2015 en el internado clínico de odontología de la Universidad de Costa Rica. *Poblac Salud Mesoam.* 2019; 16 (2): 1-18.
8. Bagan J., Sarrion G., Jimenez Y. Oral cancer: clinical features. *Oral Oncol.* 2010 Jun; 46 (6): 414-7.
9. Shulman J.D., Beach M.M., Rivera-Hidalgo F. The prevalence of oral mucosal lesions in U.S. adults: data from the Third National Health and Nutrition Examination Survey, 1988-1994. *J Am Dent Assoc.* 2004 Sep; 135 (9): 1279-86.
10. Castellanos J.L., Diaz-Guzman L. Lesions of the oral mucosa: an epidemiological study of 23785 Mexican patients. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod.* 2008 Jan; 105 (1): 79-85.
11. Barboza I., Camacho A., Gutiérrez G., Tacsan L. Epidemiología del cáncer bucal en Costa Rica en el período de 1981-1994. Seminario de graduación. Costa Rica: Facultad de Odontología, Universidad de Costa Rica; 1997.
12. Lao Gallardo W., Sobalvarro Mojica K. Egresos hospitalarios debidos a enfermedades de las glándulas salivales, CCSS, Costa Rica, 1997 al 2015. *Odontología Vital.* 2018; 28: 41-50.
13. Lao Gallardo W., Melendez Bolaños R., Herrera Jiménez A. Estudio descriptivo de cáncer bucal en los egresos hospitalarios de la Caja Costarricense de Seguro Social en los años 2001 a 2008. *Rev Cient Odontol.* 2010; 6 (2): 52-8.
14. Boza Oreamuno Y.V., López Soto A. Correlation of Clinical and Histopathological Diagnosis of Oral Mucosa Lesions at the University of Costa Rica Between 2008 and 2015. *Odovtos - International Journal of Dental Sciences.* 2020; 23 (3): 407-415.
15. Allen C.M. Examination of the oral cavity and identification of normal variants. *Dermatol Ther.* 2002; 15 (3): 174-8.
16. Hernández Carazo D., Solano Romero K., Torres Guevara E., Trejos Cisneros J.J., Fernández Chaves J.M. Determinación de la simbología más utilizada en expedientes odontológicos en Costa Rica en el año 2019 con fines de identificación de seres humanos. *Med leg Costa Rica.* 2020; 37 (1): 179-91.
17. Fernández Chaves J.M. Utilidad de expedientes odontológicos disponibles en Costa Rica en el año 2018 para la identificación de víctimas de desastres según el protocolo de INTERPOL. *Medicina Legal de Costa Rica.* 2019; 36 (1): 32-42.

18. Neville B.W., Damm D.D., Allen C.M., Chi A.C. *Oral and Maxillofacial Pathology*. 4th ed. Saunders; 2015. 928 p.
19. Thompson L.A., Chen H. *Physiology of Aging of Older Adults Systemic and Oral Health Considerations*. Clin Geriatr Med [Internet]. 2023 [cited 2023 Jul 23]; 39: 225-34. Available from: <https://doi.org/10.1016/j.cger.2023.01.009>
20. Carvalho M. de V., Iglesias D.P.P., do Nascimento G.J.F., Sobral A.P.V. Epidemiological study of 534 biopsies of oral mucosal lesions in elderly Brazilian patients. *Gerodontology*. 2011 Jun; 28 (2): 111-5.
21. Corrêa L., Frigerio M.L.M.A., Sousa S.C.O.M., Novelli M.D. Oral lesions in elderly population: a biopsy survey using 2250 histopathological records. *Gerodontology* [Internet]. 2006 Mar 1 [cited 2022 Nov 2]; 23 (1): 48-54. Available from: <https://onlinelibrary-wiley-com.ezproxy.sibdi.ucr.ac.cr/doi/full/10.1111/j.1741-2358.2006.00090.x>
22. Guedes M.M., Albuquerque R., Monteiro M., Lopes C.A., do Amaral J.B., Pacheco J.J., et al. Oral soft tissue biopsies in Oporto, Portugal: An eight year retrospective analysis. *J Clin Exp Dent*. 2015 Dec; 7 (5): e640-8.
23. Sixto-Requeijo R., Diniz-Freitas M., Torreira-Lorenzo J.C., García-García A., Gándara-Rey J.M. An analysis of oral biopsies extracted from 1995 to 2009, in an oral medicine and surgery unit in Galicia (Spain). *Med Oral Patol Oral Cir Bucal*. 2012 Jan; 17 (1): e16-22.
24. Zanin A.A., Herrera L.M., Melani R.F.H. Civil liability: characterization of the demand for lawsuits against dentists. *Braz Oral Res* [Internet]. 2016 Aug 18 [cited 2021 Sep 1]; 30 (1). Available from: <http://www.scielo.br/j/bor/a/8njNnScdJP6hmQmyZ8Qq5g/?lang=en>
25. De Lima Saintrain M.V., Holanda T.G., Bezerra T.M.M., De Almeida P.C. Prevalence of soft tissue oral lesion in elderly and its relations with deleterious habits. *Gerodontology*. 2012; 29 (2): 130-4.
26. Silva L.P., Leite R.B., Sobral A.P.V., Arruda J.A., Oliveira L.V., Noronha M.S., et al. Oral and Maxillofacial Lesions Diagnosed in Older People of a Brazilian Population: A Multicentric Study. *J Am Geriatr Soc* [Internet]. 2017 Jul 1 [cited 2022 Nov 2]; 65 (7): 1586-90. Available from: <https://onlinelibrary-wiley-com.ezproxy.sibdi.ucr.ac.cr/doi/full/10.1111/jgs.14815>
27. Vega M.J., Bedregal G.P., Jadue H.L., Delgado B.I. Equidad de género en el acceso a la atención de salud en Chile. *Rev Med Chil*. 2003 Jun; 131 (6): 669-78.
28. González Barrón S., Jiménez Corona M.E., Triana Estrada J., Ureña Ciret J.L., García Hernández J., Carrillo Rocha J. de D., et al. Recomendaciones para mejorar la práctica odontológica. *Revista CONAMED*, ISSN-e 1405-6704, Vol 8, No 1 (Enero-Marzo), 2003, págs 29-38.
29. Pinchi V., Varvara G., Pradella F., Focardi M., Donati M., Norelli G. Analysis of Professional Malpractice Claims in Implant Dentistry in Italy from Insurance Company Technical Reports, 2006 to 2010. *Int J Oral Maxillofac Implants*. 2014 Sep; 29 (5): 1177-84.
30. Consejo Dentistas de España. *La salud bucodental en España*. 2015. 1-97 p.
31. Mur G. *La Lengua en el Examen Físico*. La Homeopatía De México. 2016; 85 (704): 35-9.
32. Chiliquinga Betún E.R., Naula Vicuña C.R. Alteraciones linguales más comunes en adultos mayores. Revisión de literature Most common lingual alterations in older adults. *Review of literature. Research, Society and Development*. 2023; 12 (7): 1-8.
33. Silva L.P., Leite R.B., Sobral A.P.V., Arruda J.A., Oliveira L.V., Noronha M.S., et al. Oral and Maxillofacial Lesions Diagnosed

- in Older People of a Brazilian Population: A Multicentric Study. *J Am Geriatr Soc.* 2017 Jul 1; 65 (7): 1586-90.
34. Rivera C., Droguett D., Arenas-Márquez M.J. Oral mucosal lesions in a Chilean elderly population: A retrospective study with a systematic review from thirteen countries. *J Clin Exp Dent [Internet].* 2017 [cited 2022 Oct 15]; 9 (2): e276. Available from: /pmc/articles/PMC5303331/
 35. Ribeiro De Castro M.C., Ramos-E-Silva M. Physiologic alterations of oral mucosa due to aging: A review. *Acta Dermatovenerologica Croatica.* 2005; 13 (3): 184-8.
 36. De Lima Saintrain M.V., Holanda T.G., Bezerra T.M.M., De Almeida P.C. Prevalence of soft tissue oral lesion in elderly and its relations with deleterious habits. *Gerodontology [Internet].* 2012 Jun 1 [cited 2022 Nov 2]; 29 (2): 130-4. Available from: <https://onlinelibrary-wiley-com.ezproxy.sibdi.ucr.ac.cr/doi/full/10.1111/j.1741-2358.2011.00618.x>
 37. Carvalho M. de V., Iglesias D.P.P., do Nascimento G.J.F., Sobral A.P.V. Epidemiological study of 534 biopsies of oral mucosal lesions in elderly Brazilian patients. *Gerodontology.* 2011 Jun; 28 (2): 111-5.
 38. Lei F., Chen J.Y., Wang W.C., Lin L.M., Huang H.C., Ho K.Y., et al. Retrospective study of oral and maxillofacial lesions in older Taiwanese patients. *Gerodontology. [Internet].* 2015 [cited 2022 Nov 2]; 32 (4): 281-7. Available from: <https://onlinelibrary.wiley.com/doi/10.1111/ger.12118>
 39. Silva L.V. de O., de Arruda J.A.A., Abreu L.G., Ferreira R.C., da Silva L.P., Pelissari C., et al. Demographic and Clinicopathologic Features of Actinic Cheilitis and Lip Squamous Cell Carcinoma: a Brazilian Multicentre Study. *Head Neck Pathol.* 2020;14 (4): 899-908.
 40. Rodríguez-Blanco I., Flórez Á., Paredes-Suárez C., Rodríguez-Lojo R., González-Vilas D., Ramírez-Santos A., et al. Actinic cheilitis prevalence and risk factors: A cross-sectional, multicentre study in a population aged 45 years and over in north-west Spain. *Acta Derm Venereol.* 2018; 98 (10): 970-4.
 41. Cueto A., Martínez R., Niklander S., Deichler J., Barraza A., Esguep A. Prevalence of oral mucosal lesions in an elderly population in the city of Valparaiso, Chile. *Gerodontology.* 2013; 30 (3): 201-6.
 42. De Lima Saintrain M.V., Holanda T.G., Bezerra T.M.M., De Almeida P.C. Prevalence of soft tissue oral lesion in elderly and its relations with deleterious habits. *Gerodontology.* 2012; 29 (2): 130-4.
 43. Chi A.C., Neville B.W., Krayer J.W., Gonsalves W.C. Oral Manifestations of Systemic Disease. *American Family Physician.* 2010; 82 (11):1381-1388
 44. Hassona Y., Scully C., Abu Ghosh M., Khoury Z., Jarrar S., Sawair F. Mouth cancer awareness and beliefs among dental patients. *Int Dent J.* 2015; 65 (1): 15-21.
 45. Nagao T., Warnakulasuriya S. Screening for oral cancer: Future prospects, research and policy development for Asia. *Oral Oncol.* 2020; 105 (March): 104632.
 46. Howard Mora M., Castillo Rivas J. Cáncer bucal : conocimiento básico de la población en. *ODOVTOS-Int J Dent Sc.* 2004; (6): 94-102.
 47. Kademani D. Oral cancer. *Mayo Clin Proc.* 2007; 82 (7): 878-87.
 48. British Columbia Oral Cancer Prevention Program BCAC of DS of BColumbia. Guideline for the early detection of oral cancer in

- British Columbia 2008. *J Can Dent Assoc.* 2008; 74 (3): 245.
49. Epstein J.B., Scully C. Assessing the patient at risk for oral squamous cell carcinoma. *Spec Care Dentist.* 1997; 17 (4): 120-8.
50. World Health Organization. IARC Handbooks of Cancer Prevention Volume 19: Oral Cancer. 2020 [cited 2022 Dec 4]. IARC_Handbooks_vol_19_zoom.jpg (1500×2131). Available from: https://www.iarc.who.int/wp-content/uploads/2022/10/IARC_Handbooks_vol_19_zoom.jpg
51. Chávez E.M., Kossioni A., Fukai K. Policies Supporting Oral Health in Ageing Populations Are Needed Worldwide. *Int Dent J.* 2022; 72 (4): S27-38.