

Correlation between Personality Traits and Dimensions of Anxiety in Young Students in the State of Jalisco

Amelia Castellanos Valencia,^{1,3} Marcela Arteaga Silva,^{2,✉} Herlinda Bonilla Jaime,^{2,✉} Miguel Ángel Guevara,¹ Marisela Hernández González¹

¹ Instituto de Neurociencias, Universidad de Guadalajara, Guadalajara, Jalisco, México.

² Departamento de Biología de la Reproducción, Universidad Autónoma Metropolitana-Iztapalapa, Ciudad de México, México.

³ Doctorado en Ciencias Biológicas y de la Salud, Universidad Autónoma Metropolitana, Ciudad de México, México.

Correspondence:

Marisela Hernández González
Instituto de Neurociencias, Universidad de Guadalajara
Francisco de Quevedo 180,
Col. Arcos-Vallarta, C.P. 44130,
Guadalajara, Jalisco, México.
Phone: (52) (33) 3777-1150 ext. 33360
Email: marisela.hgonzalez@academicos.udg.mx

Received: January 12, 2023

Accepted: August 22, 2023

Citation:

Castellanos Valencia, A., Arteaga Silva, M., Bonilla Jaime, H., Guevara, M. A., & Hernández González, M. (2024). Correlation between Personality Traits and Dimensions of Anxiety in Young Students in the State of Jalisco. *Salud Mental*, 47(3), 127-135.

DOI: 10.17711/SM.0185-3325.2024.017



ABSTRACT

Introduction. Young people are highly susceptible to manifesting the various dimensions of anxiety, which have been associated with certain personality traits. **Objective.** To determine the association between personality traits (PT) and the dimensions of clinical anxiety (CA), trait anxiety (TA), and state anxiety (SA) in young students in the areas of health sciences in the state of Jalisco, Mexico. **Method.** Five hundred and twenty-one students (with an average age of 20.5 years) enrolled in psychology, nursing, and nutrition degree programs participated, 75.2% of whom were women. From April 2020 to May 2021, a cross-sectional design was used to virtually evaluate CA (Beck's inventory), TA and SA (the Spielberger inventory), and PT (Eysenck's inventory). Linear regression analysis was used to calculate correlations (R) and determination coefficients (R²) to predict the dimensions of anxiety in relation to PT. **Results.** 25.7% and 24.8% of participants scored for severe and moderate CA respectively, while 54.3% obtained high scores for TA and SA. For CA, the neuroticism PT yielded an R of .607 and an R² of .367; for TA, an R of .803 and an R² of .644; and for SA, an R of .735 and an R² of .540. **Discussion and conclusion.** The high correlation between the neuroticism PT and anxiety in young people was probably exacerbated by the restrictive conditions of the COVID-19 pandemic. In young students, the neuroticism PT is a predictor for the development of various dimensions of anxiety.

Keywords: Clinical anxiety, trait anxiety, state anxiety, neuroticism, personality traits.

RESUMEN

Introducción. Los jóvenes tienen una alta susceptibilidad a presentar manifestaciones de ansiedad en sus diferentes dimensiones, las cuales se han asociado a algunos rasgos de personalidad. **Objetivo.** Determinar la relación entre los rasgos de personalidad (RP) y las dimensiones de: ansiedad clínica (AC), ansiedad rasgo (AR) y ansiedad estado (AE) en jóvenes estudiantes de áreas de las ciencias de la salud del estado de Jalisco, México. **Método.** Participaron 521 estudiantes de psicología, enfermería y nutrición (edad promedio 20.5 años), de los cuales el 75.2% fueron mujeres. De abril 2020 a mayo 2021, mediante un diseño transversal, se evaluaron de manera virtual la AC (inventario Beck), la AR y AE (inventario Spielberger) y los RP (inventario Eysenck). Con análisis de regresión lineal se calculó la correlación (R) y el coeficiente de determinación (R²) como predictor de las dimensiones de ansiedad con los RP. **Resultados.** El 25.7% de los participantes calificó con AC grave y 24.8% con AC moderada; para la AR y AE un 54.3% obtuvo puntuaciones altas. El RP neuroticismo presentó para AC una R de .607 y R² de .367; para AR una R de .803 y R² de .644; para AE una R de .735 y R² de .540. **Discusión y conclusión.** Es probable que la alta correlación entre el RP neuroticismo y la ansiedad en los jóvenes haya sido potenciada por las condiciones restrictivas de la pandemia. El RP neuroticismo en jóvenes estudiantes, es un predictor para el desarrollo de ansiedad en sus diferentes dimensiones.

Palabras clave: Ansiedad clínica, ansiedad rasgo, ansiedad estado, neuroticismo, rasgos de personalidad.

INTRODUCTION

The anxious state is characterized by feelings of fear and worry that affect mental health and cognitive functions such as attention, memory and decision making (Bishop, 2009; Paulus & Yu, 2012), crucial to maintaining a balance between health and daily, academic, and socio-emotional skills (Simpson et al., 2010). This anxious state, common in young students (Kawakami et al., 2012), was exacerbated during the COVID-19 lockdown. Fear of the unknown and the risk of dying from infection increased anxiety levels in vulnerable young people, becoming a serious public health problem (Pérez-Cano et al., 2020; Medina-Mora, 2020).

The age of onset of the first symptoms of anxiety occurs during neurodevelopment, peaking during adolescence, a period of major age-related physical and behavioral changes. Drastic adjustments in neuronal connectivity and pruning, together with changing levels of hormones and neurotransmitters, are the basis of the functional changes associated with anxiety in late adolescence (age 24), which stabilize at the beginning of adulthood (Paus et al., 2008; Solmi et al., 2022).

Gender differences in anxiety have been described, with women having higher prevalence rates as well as more severe clinical symptoms (Asher et al., 2017). However, despite these differences in the prevalence rates of anxiety disorders, the age of onset of anxiety disorders does not differ between genders (De Lijster et al., 2017).

Anxiety is an adaptive reaction to stimuli regarded as threatening, accompanied by autonomic, neurophysiological, and psychic changes (Cattell & Scheier, 1958; Lazarus & Averill, 1972; Spielberger, 1972). According to Spielberger, in crisis situations, the immediate emotional state can be altered and modified over time depending on the biological, environmental, and social personal characteristics that explain, modulate, and maintain behavior (Spielberger, 1972).

Anxiety has various dimensions, determined by the intensity, frequency and duration of the manifestations and existence of comorbidity. It can occur as a momentary state (state-anxiety, SA) or be an intrinsic characteristic of individuals (trait-anxiety, AR). In other words, it is no longer a transitory emotional state, but a relatively stable tendency to perceive multiple situations as threatening (Spielberger, 1972; Saviola et al., 2020) and both SA and TA are evaluated using the Spielberger State-Trait Anxiety Inventory (STAI).

When these manifestations of anxiety are intense, recurrent, and also compromise social and emotional functionality and everyday activities, they are classified by the DSMV (American Psychiatric Association, 2013) as generalized anxiety disorder (GAD) or Clinical Anxiety (CA), in which the situation triggering anxiety is not congruent with the manifestations of the latter (Lee et al., 2009b).

This is evaluated using the Beck Anxiety Inventory (BAI; Beck et al., 2011).

A trait is a unit of personality, a dynamic behavioral tendency resulting from the integration of several specific habits of adjustment, expressing a characteristic mode of reaction of the individual in a given context (Allport, 1927). According to the DSM, personality traits are persistent patterns of ways of thinking, relating, and acting towards the environment and oneself that occur in a range of social settings. Various classifications of personality traits have been described (Cattell, 1943; 1945). According to Eysenck (1947), personality traits (PT) comprise four behavioral characteristics of individuals, related to ascendance (dominance and leadership), responsibility, emotional instability-anxiety, and sociability. These behavioral characteristics are grouped into two broad opposite traits: neuroticism (N), which correlates negatively with ascendance and self-esteem, and positively with emotional instability-anxiety, and extraversion (E), which correlates positively with ascendance and sociability (Eysenck, 1947; Eysenck et al., 1985). Because these behavioral tendencies have a high biological predisposition, it is difficult to tease apart the dimensions of anxiety and personality traits (Cattell & Scheier, 1958; Eysenck, 1963; Hettema et al., 2004).

Particularly in the case of young students in the areas of health sciences, the academic workload, as well as social, economic, and familial pressure during the protracted COVID-19 lockdown, constituted powerful stressors (Wathelet et al., 2020; Medina-Mora et al., 2022). Thus, the prevalence of stressful conditions, in conjunction with PTs, modulated by biological systems, result in states of anxiety in its different dimensions (Lee et al., 2009b; Caspi et al., 2010; Hansell et al., 2012). Indeed, it is known that daily exposure to various situations of emotional stress affects cognitive aspects of attention, interpretation and memory that facilitate the onset and maintenance of anxiety (Clarke et al., 2008), resulting in the poor academic performance of individuals, especially those of school age (Vilaplana-Pérez et al., 2021). However, it is unclear to what extent personality type determines the likelihood of presenting with anxiety. Establishing associations between PT and the various dimensions of anxiety in young students will therefore make it possible to enhance our knowledge of the interaction between personality and anxiety presented by youth and how this interaction could influence or impact emotional problems, as well as the cognitive and behavioral alterations prevailing in this sector of the population (Lee et al., 2009a).

Given the difficulty of teasing apart personality traits from anxiety dimensions, the hypothesis of this paper was that the neuroticism personality trait would present different degrees of correlation and prediction with anxiety dimensions depending on sex, age, and academic performance in young students from the state of Jalisco during the pandemic.

It was also hypothesized that the detection of personality traits predicting anxiety dimensions and dysfunctional behavioral patterns in young students would allow the use of early intervention strategies.

METHOD

Study design

This was a cross-sectional study with non-probabilistic sampling, conducted in the state of Jalisco, Mexico in May 2021.

Participants

The first step was to invite all the students enrolled in the psychology, nursing, and nutrition degree programs (young women and men ages 18-30) at two university centers: the University of Guadalajara, Ciénega de Ocotlán campus (CUCI UDG) and the University of Specialties Health Sciences campus (UNE). Students were informed that this was a research study, which complied with ethical guidelines, and had been previously authorized by the Ethics in Research Commission of the Institute of Neurosciences, based on NIH guidelines. All study participants voluntarily answered the survey.

Procedure

Information was provided through a link sent via email and social networks. Students were invited to participate on the understanding they could drop out of the survey if they so wished. In addition, they were informed that the data collected in this study would be held anonymously and be for the exclusive use of the laboratory.

Participants were told that this was an individual self-assessment survey, and that there were no right or wrong answers. They were asked not to spend a lot of time answering each question and to attempt to provide the answer that best indicated how they described themselves.

Google Forms was used for the virtual administration of the three anxiety scales, as well as the personality traits scale. Once participants had opened the link, they were able to access the information section, as well as the questionnaires and inventories required to ensure that they met the inclusion criteria, which began with a sociodemographic questionnaire with questions on their age, sex, and grades. Immediately afterwards, instructions for answering each inventory appeared on the screen. The instructions were as follows: “Read every sentence and fill in the numbered circle indicating how you feel usually or right now” according to the question. First, the 21-item Beck Clinical Anxiety Inventory (BAI) appeared. This was followed by the State-

Trait Anxiety Inventory (STAI), comprising twenty items for TA and 20 for SA, and finally, a screen with the 24-item personality questionnaire. Once students had completed the survey, a thank you message appeared on the screen, indicating the end of the survey. The link closed and the results were automatically saved.

Scales and questionnaires

The Beck Anxiety Inventory (BAI) is a scale measuring the severity of clinical anxiety in adults and adolescents. The twenty-one items in the BAI contain the DSMV symptomatic criteria for the diagnosis of clinical anxiety, (including anxiety or panic attacks), for the last week including the day of the survey (Beck et al., 2011). In this paper, scores of 0-7 were described as “minimal CA.” Scores of 8-15 were termed “mild CA.” Scores of 16-25 were classified as “Moderate CA” and lastly, scores of 26-63 were described as “Severe CA” (Beck et al., 2011).

The State-Trait Anxiety Inventory (STAI) adapted to the Mexican population (Spielberger & Díaz-Guerrero, 2002) was used to measure the two dimensions of anxiety. The STAI Inventory has a section to assess trait anxiety (TA) consisting of twenty affirmative statements describing how the participant usually feels, whereas in the section to assess state anxiety (SA), the twenty expressions indicate how the participant feels at that moment (right now). Participants who obtained a score over 44 were considered to be in the TA or high SA group, while those who obtained a score below 30 were considered to be in the TA or low SA group (Spielberger & Díaz-Guerrero, 2002).

The present study was conducted in two phases.

In the first phase of the study, personality traits were assessed using the revised Eysenck personality questionnaire, abbreviated form (EPQR-A; Francis et al., 1992). The latter consists of twenty-four items divided into four dimensions of six items per subscale: extraversion (E), including aspects of positive emotionality, sociability, spontaneity, vitality and externalization; neuroticism (N), including traits such as negative emotionality, anxiety, sensitivity, worry, and self-consciousness; psychoticism (P), encompassing aspects of cruelty, impulsivity, low socialization, non-conformity, irresponsibility and schizoidism, and finally, dissimulation or lying (L), referring to the propensity to conceal or lie (Eysenck et al., 1985). A Likert-type scale response format was used (Muñiz et al., 2005), while the dispersion for each subscale varied from a minimum of four to a maximum of twenty-four points with a median of twelve. In this paper, direct scores were obtained from each subscale, without T transformation.

In the second phase of the study, only one personality trait, neuroticism (N), was evaluated using the revised Eysenck personality questionnaire, with twenty-four items corresponding to the complete EPQR-N neuroticism sub-

scale, with dichotomous responses (Yes/No); and a maximum score of twenty-four (Eysenck et al., 1985).

To determine which personality trait contributed or explained the most in each of the dimensions of anxiety in the study participants, a sample of 185 students classified into different groups by age, sex and school grade were included in the first phase. CA, TA, SA, and Personality Trait scores were determined with the abbreviated scale (EPQR-A) for N, P, E and L using the scales and questionnaires mentioned earlier.

Due to their low contribution and to prevent oversaturation of the predictive model, to obtain a simplified model with the contribution of valid and replicable predictions, the elimination of PTs that contributed little to predicting anxiety dimensions was proposed. For each predictor introduced into the model, adjusted R^2 , there was a penalty since it depended on the degrees of freedom.

Data analysis

The reliability of the scales was established with Cronbach's alpha (Cronbach, 1951). The raw total scores of Beck's Anxiety Inventory, Spielberger's State-Trait Anxiety Inventory (STAI) and the Eysenck Personality Profile Inventory were described and compared between groups (two age groups 18-24 and 25-30 years), two groups by sex (men and women) and three groups by grades (70-80, 81-90 and 91-100) as categorical independent variables. The difference between the theoretical and observed cumulative distribution of the variables described was calculated with the Kolmogorov-Smirnov (KS) test, to determine which statistic to use to compare the differences in means or medians and standard deviation (SD) or ranges. When there were two independent groups and normal distribution according to KS, the parametric T-test was used. If the distribution was not normal according to KS, non-parametric, Mann Whitney U test was used. For three groups,

according to their distribution, one-way ANOVA was used, followed by Tukey's test. Values of $p < .05$ were considered significant.

Using multiple linear regression analysis and statistics to determine whether there was collinearity with the Durbin-Watson test, the correlation (R) and the coefficient of determination (R^2) were calculated as a predictor of anxiety dimensions with all PTs, in addition to sociodemographic variables. IBM SPSS Statistics 25 was used.

The sample size was calculated using the latest version of G*Power (v 3.1.9.7; Heinrich-Heine-Universität Düsseldorf, Düsseldorf, Germany). With a power ($1-\beta$) of .80, and a significance level (α) of .05, for a bivariate Pearson linear correlation analysis (with an a priori calculation from the worst-case scenario of a low correlation of 0.2), a sample size of 193 subjects would be required. However, based on previous studies in which a correlation of neuroticism with trait anxiety of .848 and of .70 with state anxiety was obtained (Eid et al., 2022), the necessary sample size was eight to thirteen subjects.

Ethical considerations

None of the procedures in this non-invasive research posed a risk to the physical, emotional, or moral integrity of the participants. The research therefore qualifies as minimal risk in accordance with Article 17 of the Regulations of the General Health Law on Health Research. In addition, the guidelines specified in the 1975 Declaration of Helsinki and its amendments were followed, as well as current international codes and standards for good practices in research with humans, and the Official Mexican Standard NOM-012SSA3-2012, which establishes the criteria for undertaking research projects for health in human beings.

This project was approved by the Ethics Committee of the Institute of Neurosciences of the University of Guadalajara, with registration number ET022021-314.

Table 1
Raw Scores of Anxiety Dimensions and Personality Traits in the Various Groups of Young Students

	<i>n</i> = 185	CA	TA	SA	N	E	P	L
Age								
18-24	168 (90.81%)	17.5 ± 11 ^a	46.5 ± 10 ^a	46 ± 10	13.8 ± 4.2 ^a	14.1 ± 2.6 ^a	13.4 ± 2.3	10.9 ± 2.1
25-30	17 (9.19%)	11.1 ± 8	39.9 ± 9	41.2 ± 9	10.5 ± 3.6	12.1 ± 3.2	13 ± 2.3	10.5 ± 1.7
Sex								
Female	138 (74.6%)	17 ± 8 ^b	47 ± 8 ^b	47 ± 9 ^b	14 ± 3 ^b	13 ± 3	13 ± 1	11 ± 1
Male	47 (25.4%)	13 ± 7	41 ± 10	41.6 ± 8	12 ± 3	14 ± 2	14 ± 2	11 ± 1
Grades								
70-80	7 (4%)	19.1 ± 11	51.4 ± 8.3	47.2 ± 9.8	14 ± 4.8	14.2 ± 2.8	13.8 ± 3.4	10.2 ± 1.3
81-90	81 (46.6%)	17.5 ± 12	45.5 ± 9.9	46.4 ± 10.9	13.4 ± 4.3	13 ± 2.6	13.4 ± 2.4	10.7 ± 2
91-100	86 (49.4%)	16.7 ± 12	46.2 ± 10	44.8 ± 10.8	13.6 ± 4.3	13.2 ± 3	14.2 ± 2.1	11.2 ± 2

Note: ^a = Mean ± DS, ^b = Median ± CI range of the raw scores of Clinical anxiety (CA), Trait anxiety (TA), State anxiety (SA) and EPQR-A personality trait: Neuroticism (N), Extraversion (E), Psychoticism (P) and Lying (L), displayed by young people in groups by age, sex and grade. *n* = 185; the significant differences, $p < .05$, are in regard to the same group. Based on tests from ^a T-test, ^b Mann Whitney, ^c ANOVA.

Table 2
Predictive Model of Personality Traits for Anxiety Dimensions (n = 185)

	CA			TA			SA		
	R	Adjusted R ²	Sig. Change in F	R	Adjusted R ²	Sig. Change in F	R	Adjusted R ²	Sig. Change in F
N	.686	.468	< .001	.782	.609	< .001	.759	.574	< .001
E	.020	-.005	.782	.017	-.005	.814	.091	.003	.217
P	.146	.016	.048	.139	.014	.059	.198	.034	.007
L	.055	-.002	.459	.024	-.005	.743	.063	-.002	.398
Age	.222	.044	.002	.211	.039	.004	.167	.023	.023
Sex	.196	.033	.007	.191	.031	.009	.183	.028	.013
Grade	.024	-.005	.757	.039	-.004	.610	.039	-.004	.611

Note: Linear regression analysis. Predictors: personality traits, Neuroticism (N), Extraversion (E), Psychoticism (P), Lying (M), age, sex, and grade. Dependent variable: Clinical anxiety (CA), Trait anxiety (TA), State anxiety (SA). n = 185, R = Pearson correlation, R² = Coefficient of determination, Sig = statistical significance.

RESULTS

Phase 1

Subjects who did not fall within the age range of 18 to 30 or failed to complete the survey were eliminated. Of the 185 participants who completed the entire survey at various university centers in the state of Jalisco, Mexico, with an average age of 21 (± 2.6) years, 74% were women; It was found that those under 24 and women had a higher total score for CA, as well as TA and SA. Students with lower grades had higher total CA and SA scores. Participants under 24 and women also displayed higher PTs for neuroticism (Table 1).

The reliability found in the Phase 1 sample of 185 participants or Cronbach's alpha coefficient was .923. for the BAI CA scale, .888 for the STAI TA, .913 for the STAI SA and .823 for the EPQR-A personality traits.

Subsequently, to determine and predict how much each of the personality traits contributed to the anxiety dimensions, a multiple linear regression analysis was conducted,

with statistics to determine whether there was collinearity. The correlation and determination coefficient were determined, using the four personality traits, age, sex, and grades, as predictor variables for each of the dimensions of CA, TA, and SA as dependent variables. Values of *p* < .05 were considered significant. All of the above was analyzed using SPSS Statistics 25.

The predictor values for CA, TA, and SA are described in Table 2. As can be seen, the neuroticism personality trait achieved the highest statistically significant predictor value, indicating that this personality trait contributes most to the presence of CA, TA, and SA. Psychoticism, age, and sex only contributed minimally while the other traits provided little or no predictive value.

Phase 2

According to the results of Phase 1, the neuroticism personality trait is the one most strongly associated with the various dimensions of anxiety. Thus, based on the results

Table 3
Raw Scores of Anxiety Dimensions and Personality Traits in the Various Groups of Young Students

	n = 521	CA	TA	SA	N
Age					
18-24	488 (93%)	18.3 ± 12 ^a	46.6 ± 10 ^a	46.2 ± 12	13.7 ± 5.8 ^a
25-30	33 (7%)	13.6 ± 11	42.7 ± 11	44.3 ± 13	11.2 ± 6.5
Sex					
Female	392 (75.2%)	17 ± 9 ^b	47 ± 8 ^b	47 ± 8 ^b	15 ± 5 ^b
Male	129 (24.8%)	14 ± 9	43 ± 9	42 ± 10	12 ± 6
Grade					
70-80	21 (4%)	21.4 ± 14	52.8 ± 10 ^c	52.8 ± 13 ^c	14.1 ± 5
81-90	115 (24%)	18.3 ± 11	46.6 ± 10 ^c	48 ± 12	13.3 ± 6
91-100	346 (72%)	17.6 ± 11	45.9 ± 10.8 ^c	45.1 ± 11 ^c	13.5 ± 5.8

Note: ^a = Mean ± SD, ^b = Median ± Range CI of the raw scores of Clinical anxiety (CA), Trait anxiety (TA), State anxiety (SA) and the Neuroticism personality trait (N), presented by the young people in groups by age, sex and grade, n = 521. The significant differences, *p* < .05, are in regard to the same group. Based on tests from ^a T-test, ^b Mann Whitney, ^c ANOVA post hoc Tukey.

Table 4
Predictive Factors: neuroticism personality traits, age, sex and grade for dimensions of anxiety

	CA			TA			SA		
	R	Adjusted R ²	Sig. Change in F	R	Adjusted R ²	Sig. Change in F	R	Adjusted R ²	Sig. Change in F
N	.592	.349	< .001	.803	.644	< .001	.735	.540	< .001
Age	.097	.007	.027	.086	.005	.050	.039	.002	.375
Sex	.102	.008	.020	.195	.036	< .001	.143	.018	.001
Grade	.057	.001	.215	.105	.009	.021	.153	.022	.001

Note: Linear regression analysis. Predictors: personality traits, Neuroticism (N), age, sex, and school grade. Dependent variables Clinical anxiety (CA), Trait anxiety (TA), State anxiety (SA) $n = 521$, R = Pearson correlation, R²: Coefficient of determination, Sig: statistical significance.

of Phase 1, a second phase of sampling and analysis was conducted, with a sample of 521 students also classified into different groups by age, sex, and grades. In this sample, the CA, TA, SA, and Personality Trait scores were also determined, but only with the complete neuroticism subscale (EPQR-N; Eysenck et al., 1985) to be able to associate and predict how much neuroticism contributes to each of the dimensions of anxiety. To this end, a Pearson linear regression analysis and coefficient of determination (adjusted R²) were applied, to show how much neuroticism predicts the dimensions of anxiety.

As in phase 1, a demographic description of the participants was made, shown in Table 3. Subjects who did not fall within the age range of 18 to 30 or failed to complete the survey were eliminated. In the end, a total of 521 students from various university centers in the state of Jalisco, Mexico participated. Enrolled in the areas of Health Sciences, students had an average age of 20.5 (\pm 2.3) years, and 75.2% were women and 24.8% men. It was found that those under 24 and women had a higher total score for CA, as well as for TA, while only women had SA. Students with lower grades had higher total CA and SA scores. Participants under 24 and women also displayed higher PTs for neuroticism (Table 3).

As with the data from Phase 1, a regression analysis was conducted to determine the correlation and determination coefficient, using only the data from the complete neuroticism scale (EPQR-N), age, sex, and grades, as predictor

variables for each of the dimensions of CA, TA, and SA as dependent variables. As can be seen in Table 4, demographic variables (age, sex, and school grade) contributed little to predicting the dimensions of anxiety.

In addition, a regression analysis was conducted to determine the final predictive model using only data from the complete neuroticism scale (EPQR-N) as a predictor variable and the anxiety dimensions as dependent variables: CA, TA, and SA.

It was found that the neuroticism personality trait contributed to the various dimensions of anxiety, in varying proportions: .367 for CA, .644 for TA and .540 for SA. Likewise, it was found that neuroticism predicts all dimensions of anxiety with a coefficient of determination (adjusted R²) of .655 (Table 5).

DISCUSSION AND CONCLUSION

An individual's personality trait is derived from the interaction of a series of genetic and environmental factors, characterizing the differences found in each individual (Vukasović & Bratko, 2015; Gross & Hen, 2004). This paper found that the biological personality trait characterized by negative emotionality, neuroticism, contributed most to the presence of anxiety in its various dimensions with a proportion of over 60% in TA, surpassing other factors such as sex, age, grades, or other PT showing predictive independence.

Table 5
Predictive model of neuroticism personality trait, for anxiety dimensions (n = 521)

	R	R ²	Adjusted R ²	Change in F	Sig. Change in F	Durbin-Watson
CA	.607	.368	.367	302.215	< .001	1.835
TA	.803	.645	.644	942.398	< .001	1.78
SA	.735	.55	.54	61.412	< .001	1.918
All anxiety dimensions	.81	.657	.655	329.366	< .001	1.875

Note: Linear regression analysis. Predictor = Neuroticism personality trait (N), Independent variable = Clinical anxiety (CA), Trait anxiety (TA), State anxiety (SA) and all dimensions of anxiety together. $n = 521$; R = Pearson correlation; R² = Coefficient of determination; Sig = statistical significance.

From the 1940s onwards, Eysenck made significant contributions to the description of human PTs (Eysenck, 1944; 1947; 1963; Eysenck et al., 1985), a classification that has continued to be used until the present (Eysenck, 2016). His contribution laid the foundations for determining that individual differences in personality traits are associated with cognitive processes and neural networks (Eysenck, 1997; Saviola et al., 2020). These individual differences in personality traits determine the way situations are perceived as threatening and the anxiety response. Knowing the personality trait of everyone would therefore make it possible to predict their anxiety reaction to a threat.

The results of this work show that the neuroticism PT is most strongly associated with the presence of the three dimensions of anxiety in the students in health areas in our sample. These results coincide with what has been reported by Bienvenu & Stein (2003) and other authors (Bienvenu & Stein, 2003; Brandes & Bienvenu, 2006; Kotov et al., 2007), who found that certain PTs are associated with the presence of anxiety, neuroticism being the one most associated with anxiety disorder; However, they did not distinguish between the three dimensions of anxiety, which is the original contribution of the present study.

Other research has reported that the biological personality trait neuroticism is genetically associated with anxiety (Hettema et al., 2004; Barlow et al., 2014; Zugliani et al., 2017). Our results show that the greatest correlation and predictive value of neuroticism is with the presence of TA (Clarke et al., 2008; Uliaszek et al., 2009; Zugliani et al., 2017). The high correlation obtained in our sample was probably due to the fact that neuroticism and TA share common features, as has been suggested by other authors (Uliaszek et al., 2009).

Likewise, the results of other reports concur with ours. These reports observe an even smaller contribution than the one we found in this paper of neuroticism to the prediction of the manifestations of clinical anxiety or panic, which can alter physical homeostasis. If left uncontrolled, its intense, recurrent manifestations can result in severe generalized anxiety disorder (Jorm et al., 2000; Brown & Naragon-Gainey, 2013; Zugliani et al., 2017; Pérez-Mengual et al., 2021).

In general terms, women were found to have a greater prevalence and severity of manifestations of dimensions of anxiety, coinciding with other studies in which sex differences in personality and anxiety traits are reported in all its dimensions (Hettema et al., 2005; Gottschalk & Domschke, 2017).

Our data differ from those found by Eid et al. (2022) who reported differences between TA and SA with values similar to those obtained in our group of young people. Neuroticism was found to contribute to a lesser extent to the momentary reaction of SA in participants' self-reports.

One of the most striking results of the present study is that overall, a higher prevalence of anxiety was found than that reported by other authors (Wathelet et al., 2020; Vahra-

tian et al., 2021; Santomauro et al., 2021; Medina-Mora et al., 2022). This difference is probably due to the restrictive conditions of the pandemic experienced between 2020 and 2021. The lack of information on the conditions and aspects of how each participant experienced lockdown, and only having answers obtained virtually, making it impossible to control the way the surveys were administered (ideally individually and in a setting with no distractions), meant that the data compilation was out of our hands.

For over twenty years, mental health has been a priority in the surveillance and treatment system for the WHO, due to the fact that a considerable increase in emotive-emotional diseases has been reported (Twenge, 2000). And in Mexico, the National Survey of Psychiatric Epidemiology (ENEP) has reported an over two-digit increase in the prevalence of anxiety disorders (Medina-Mora et al., 2003; Benjet et al., 2016; Medina-Mora et al., 2022), underlining the importance of establishing more controlled monitoring mechanisms nationwide.

Although the results obtained in the present study only constitute a sample of the high correlation between the neuroticism PT and the presence of anxiety in the youth population of the state of Jalisco, these data provide evidence of the severity of this health problem that probably occurs at the national level, particularly among young students in the area of health sciences. There is therefore an urgent need for the health system and the family nucleus to design more strategies to support and prevent anxiety especially in subjects with high biological vulnerability due to their neuroticism personality trait, which, together with external factors such as the pandemic and its consequences, increased the prevalence of the dimensions of anxiety among young people in the state of Jalisco.

Funding

Partly funded by the Federal Neurosciences Grant number 256063.

Conflict of interests

The authors declare that they have no conflicts of interest.

REFERENCES

- Allport, G. (1927). Concepts of trait and personality. *Psychological Bulletin*, 24(5), 284-293. doi: 10.1037/h0073629
- American Psychiatric Association. (2013). *Diagnostic and statistical manual of mental disorders: DSM-5™* (Fifth Ed). Washington DC: American Psychiatric Association.
- Asher, M., Asnaani, A., & Aderka, I. (2017). Gender differences in social anxiety disorder: A review. *Clinical Psychology Review*, 56, 1-12. doi: 10.1016/j.cpr.2017.05.004
- Barlow, D., Ellard, K., Sauer-Zavala, S., Bullis, J., & Carl, J. (2014). The Origins of Neuroticism. *Perspectives on Psychological Science*, 9(5), 481-496. doi: 10.1177/1745691614544528
- Beck, A., Steer, R., & Sanz, J. (2011). *Beck Anxiety Inventory BAI*. Madrid: Pearson.
- Benjet, C., Borges, G., Méndez, E., Albor, Y., Casanova, L., Orozco, R., Curiel, T., Fleiz, C., & Medina Mora, M. (2016). Eight-year incidence of psychiatric

- disorders and service use from adolescence to early adulthood: longitudinal follow-up of the Mexican Adolescent Mental Health Survey. *European Child & Adolescent Psychiatry*, 25, 163-173. doi: 10.1007/s00787-015-0721-5
- Bienvenu, O., & Stein, M. (2003). Personality and anxiety disorders: A review. *Journal of Personality Disorders*, 17(2), 139-151. doi: 10.1521/pedi.17.2.139.23991
- Bishop, S. (2009). Trait anxiety and impoverished prefrontal control of attention. *Nature Neuroscience*, 12(1), 92-98. doi: 10.1038/nn.2242
- Brandes, M., & Bienvenu, O. (2006). Personality and anxiety disorders. *Current Psychiatry Reports*, 8(4), 263-269. doi: 10.1007/s11920-006-0061-8
- Brown, T., & Naragon-Gainey, K. (2013). Evaluation of the Unique and Specific Contributions of Dimensions of the Triple Vulnerability Model to the Prediction of DSM-IV Anxiety and Mood Disorder Constructs. *Behavior Therapy*, 44(2), 277-292. doi: 10.1016/j.beth.2012.11.002
- Caspi, A., Hariri, A., Holmes, A., Uher, R., & Moffitt, T. (2010). Genetic sensitivity to the environment: the case of the serotonin transporter gene and its implications for studying complex diseases and traits. *American Journal of Psychiatry*, 167(5), 509-527. doi: 10.1176/appi.ajp.2010.09101452
- Cattell, R. (1943). The description of personality: basic traits resolved into clusters. *The Journal of Abnormal and Social Psychology*, 38(4), 476-506. doi: 10.1037/h0054116
- Cattell, R. (1945). The diagnosis and classification of neurotic states; a reinterpretation of Eysenck's factors. *Journal of Nervous and Mental Disease*, 102(6), 576-589. doi: 10.1097/00005053-194512000-00005
- Cattell, R., & Scheier, I. (1958). The Nature of Anxiety: A Review of Thirteen Multivariate Analyses Comprising 814 Variables. *Psychological Reports*, 4(3), 351-388. doi: 10.2466/pr0.1958.4.3.351
- Clarke, P., MacLeod, C., & Shirazee, N. (2008). Prepared for the worst: Readiness to acquire threat bias and susceptibility to elevate trait anxiety. *Emotion*, 8(1), 47-57. doi: 10.1037/1528-3542.8.1.47
- Cronbach, L. (1951). Coefficient alpha and the internal structure of tests. *Psychometrika*, 16(3), 297-334. doi: 10.1007/BF02310555
- De Lijster, J., Dierckx, B., Utens, E., Verhulst, F., Zieldorff, C., Dieleman, G., & Legerstee, J. (2017). The Age of Onset of Anxiety Disorders. *Canadian Journal of Psychiatry*, 62(4), 237-246. doi: 10.1177/0706743716640757
- Eid, C., Hamilton, C., & Greer, J. (2022). Untangling the tingle: Investigating the association between the Autonomous Sensory Meridian Response (ASMR), neuroticism, and trait & state anxiety. *PLoS One*, 17(2), e0262668. doi: 10.1371/journal.pone.0262668
- Eysenck, H. (1944). Types of personality: a factorial study of seven hundred neurotics. *Journal of Mental Science*, 90(381), 851-861. doi: 10.1192/bjp.90.381.851
- Eysenck, H. (1947). *Dimensions of Personality*. London: Routledge & Kegan Paul.
- Eysenck, H. (1963). Biological Basis of Personality. *Nature*, 199, 1031-1034. doi: 10.1038/1991031a0
- Eysenck, M. (1997). *Anxiety and cognition: A unified theory*. Hove, UK: Psychology Press. doi: 10.4324/9781315804606
- Eysenck, M. (2016). Hans Eysenck: A research evaluation. *Personality and Individual Differences*, 103, 209-219. doi: 10.1016/j.paid.2016.04.039
- Eysenck, S., Eysenck, H., & Barrett, P. (1985). A revised version of the Psychoticism Scale. *Personality and Individual Differences*, 6(1), 21-29. doi: 10.1016/0191-8869(85)90026-1
- Francis, L., Brown, L., & Philipchalk, R. (1992). The development of an abbreviated form of the revised Eysenck personality questionnaire (EPQR-A): Its use among students in England, Canada, the U.S.A. and Australia. *Personality and Individual Differences*, 13(4), 443-449. doi: 10.1016/0191-8869(92)90073-X
- Gottschalk, M., & Domschke, K. (2017). Genetics of generalized anxiety disorder and related traits. *Dialogues in Clinical Neuroscience*, 19(2), 159-168. doi: 10.31887/DCNS.2017.19.2/kdomschke
- Gross, C., & Hen, R. (2004). Genetic and Environmental Factors Interact To Influence Anxiety. *Neurotoxicity Research*, 6(6), 493-501. doi: 10.1007/BF03033286
- Hansell, N., Wright, M., Medland, S., Davenport, T., Wray, N., Martin, N., & Hickie, I. (2012). Genetic co-morbidity between neuroticism, anxiety/depression and somatic distress in a population sample of adolescent and young adult twins. *Psychological Medicine*, 42(6), 1249-1260. doi: 10.1017/S0033291711002431
- Hettema, J., Prescott, C., & Kendler, K. (2004). Genetic and environmental sources of covariation between generalized anxiety disorder and neuroticism. *American Journal of Psychiatry*, 161(9), 1581-1587. doi: 10.1176/appi.ajp.161.9.1581
- Hettema, J., Prescott, C., Myers, J., Neale, M., & Kendler, K. (2005). The Structure of Genetic and Environmental Risk Factors for Anxiety Disorders in Men and Women. *Archives of General Psychiatry*, 62(2), 182-189. doi: 10.1001/archpsyc.62.2.182
- Jorm, A., Christensen, H., Henderson, A., Jacomb, P., Korten, A., & Rodgers, B. (2000). Predicting anxiety and depression from personality: Is there a synergistic effect of neuroticism and extraversion? *Journal of Abnormal Psychology*, 109(1), 145-149. doi: 10.1037/0021-843X.109.1.145
- Kawakami, N., Abdulghani, E., Alonso, J., Bromet, E. J., Bruffaerts, R., Caldas-de-Almeida, J., Chiu, W. T., de Girolamo, G., de Graaf, R., Fayyad, J., Ferry, F., Florescu, S., Gureje, O., Hu, C., Lakoma, M., Leblanc, W., Lee, S., Levinson, D., Malhotra, S., ... Kessler, R. (2012). Early-life mental disorders and adult household income in the World Mental Health Surveys. *Biological Psychiatry*, 72(3), 228-237. doi: 10.1016/j.biopsych.2012.03.009
- Kotov, R., Watson, D., Robles, J., & Schmidt, N. (2007). Personality traits and anxiety symptoms: the multilevel trait predictor model. *Behaviour Research and Therapy*, 45(7), 1485-1503. doi: 10.1016/j.brat.2006.11.011
- Lazarus, R., & Averill, J. (1972). Emotion and cognition: with special reference to anxiety. In C. Spielberger. *Anxiety Current Trends in Theory and Research* (pp. 241-283). New York: Academic Press.
- Lee, S., Tsang, A., Breslau, J., Aguilar-Gaxiola, S., Angermeyer, M., Borges, G., Bromet, E., Bruffaerts, R., de Girolamo, G., Fayyad, J., Gureje, O., Haro, J., Kawakami, N., Levinson, D., Oakley Browne, M., Ormel, J., Posada-Villa, J., Williams, D., & Kessler, R. (2009a). Mental disorders and termination of education in high-income and low- and middle-income countries: epidemiological study. *British Journal of Psychiatry*, 194(5), 411-417. doi: 10.1192/bjp.bp.108.054841
- Lee, S., Tsang, A., Ruscio, A., Haro, J., Stein, D., Alonso, J., Angermeyer, M. C., Bromet, E. J., Demyttenaere, K., de Girolamo, G., de Graaf, R., Gureje, O., Iwata, N., Karam, E. G., Lepine, J.-P., Levinson, D., Medina-Mora, M. E., Oakley Browne, M. A., Posada-Villa, J., & Kessler, R. (2009b). Implications of modifying the duration requirement of generalized anxiety disorder in developed and developing countries. *Psychological Medicine*, 39(7), 1163-1176. doi: 10.1017/S0033291708004807
- Medina-Mora, M. (2020). COVID-19 and mental health: Challenges and opportunities. *Salud Mental*, 43(6), 241-242. doi: 10.17711/sm.0185-3325.2020.033
- Medina-Mora, M., Borges, G., Lara, C., Benjet, C., Blanco, J., Fleiz, C., Villatoro, J., Rojas, E., Zambrano, J., Casanova, L., & Aguilar, G. (2003). Prevalence of mental disorders and use of services: Results from the Mexican National Survey of Psychiatric Epidemiology. *Salud Mental*, 26(4), 1-16.
- Medina-Mora, M., Guerrero, B., & Cortés, J. (2022). ¿Cómo impactó a la salud mental de los estudiantes mexicanos la retracción de la economía resultado del largo periodo del confinamiento por COVID-19? *Boletín COVID-19 Salud Pública*, 3(28), 3-6.
- Muñiz, J., García-Cueto, E., & Lozano, L. (2005). Item format and the psychometric properties of the Eysenck Personality Questionnaire. *Personality and Individual Differences*, 38(1), 61-69. doi: 10.1016/j.paid.2004.03.021
- Paulus, M., & Yu, A. (2012). Emotion and decision-making: affect-driven belief systems in anxiety and depression. *Trends in Cognitive Sciences*, 16(9), 476-483. doi: 10.1016/j.tics.2012.07.009
- Paus, T., Keshavan, M., & Giedd, J. (2008). Why do many psychiatric disorders emerge during adolescence? *Nature Reviews Neuroscience*, 9, 947-957. doi: 10.1038/nrn2513
- Pérez-Cano, H., Moreno-Murguía, M., Morales-López, O., Crow-Buchanan, O., English, J., Lozano-Alcázar, J., & Somilleda-Ventura, S. (2020). Anxiety, depression, and stress in response to the coronavirus disease-19 pandemic. *Cirugía y Cirujanos*, 88(5), 562-568. doi: 10.24875/ciru.20000561
- Pérez-Mengual, N., Aragonés-Barbera, I., Moret-Tatay, C., & Moliner-Albero, A. (2021). The Relationship of Fear of Death Between Neuroticism and Anxiety During the Covid-19 Pandemic. *Frontiers in Psychiatry*, 20(12), 648498. doi: 10.3389/fpsy.2021.648498
- Santomauro, D., Mantilla Herrera, A., Shadid, J., Zheng, P., Ashbaugh, C., Pigott, D., Abbafati, C., Adolph, C., Amlag, J., Aravkin, A., Bang-Jensen, B., Bertolacci,

- G., Bloom, S., Castellano, R., Castro, E., Chakrabarti, S., Chattopadhyay, J., Cogen, R., Collins, J., ... Ferrari, A. (2021). Global prevalence and burden of depressive and anxiety disorders in 204 countries and territories in 2020 due to the COVID-19 pandemic. *Lancet*, *398*(10312), 1700-1712. doi: 10.1016/S0140-6736(21)02143-7
- Saviola, F., Pappaianni, E., Monti, A., Grecucci, A., Jovicich, J., & De Pisapia, N. (2020). Trait and state anxiety are mapped differently in the human brain. *Scientific Reports*, *10*(1), 11112. doi: 10.1038/s41598-020-68008-z
- Simpson, H., Neria, Y., Lewis-Fernandez, R., & Schneier, F. (2010). *Anxiety disorders – theory, research and clinical*. Cambridge: Cambridge University Press.
- Solmi, M., Radua, J., Olivola, M., Croce, E., Soardo, L., Salazar de Pablo, G., Shin, J. I., Kirkbride, J. B., Jones, P., Kim, J. H., Kim, J. Y., Carvalho, A. F., Seeman, M. V., Correll, C. U., & Fusar-Poli, P. (2022). Age at onset of mental disorders worldwide: large-scale metaanalysis of 192 epidemiological studies. *Molecular Psychiatry*, *27*(1), 281-295. doi: 10.1038/s41380-021-01161-7
- Spielberger, C. (1972). Anxiety as an emotional state. In A. Press (Ed). *Anxiety Behavior* (pp. 23-49). New York: C. D. Spielberger.
- Spielberger, C., & Díaz-Guerrero, R. (2002). *IDARE Inventario de ansiedad: Rasgo-Estado*. México: El Manual Moderno.
- Twenge, J. (2000). The Age of Anxiety? Birth Cohort Change in Anxiety and Neuroticism, 1952-1993. *Journal of Personality and Social Psychology*, *79*(6), 1007-1021. doi: 10.1037//0022-3514.79.6.1007
- Uliaszek, A., Hauner, K., Zinbarg, R., Craske, M., Mineka, S., Griffith, J., & Rose, R. (2009). An Examination of Content Overlap and Disorder-Specific Predictions in the Associations of Neuroticism with Anxiety and Depression. *Journal of Research in Personality*, *43*(5), 785-794. doi: 10.1016/j.jrp.2009.05.009
- Vahratian, A., Blumberg, S., Terlizzi, E., & Schiller, J. (2021). Symptoms of Anxiety or Depressive Disorder and Use of Mental Health Care Among Adults During the COVID-19 Pandemic — United States, August 2020–February 2021. *MMWR. Morbidity and Mortality Weekly Report*, *70*(13), 490-494. doi: 10.15585/mmwr.mm7013e2
- Vilaplana-Pérez, A., Pérez-Vigil, A., Sidorchuk, A., Brander, G., Isomura, K., Hesselmark, E., Kuja-Halkola, R., Larsson, H., Mataix-Cols, D., & Fernández de la Cruz, L. (2021). Much more than just shyness: the impact of social anxiety disorder on educational performance across the lifespan. *Psychological Medicine*, *51*(5), 861-869. doi: 10.1017/S0033291719003908
- Vukasović, T., & Bratko, D. (2015). Heritability of personality: A meta-analysis of behavior genetic studies. *Psychological Bulletin*, *141*(4), 769-785. doi: 10.1037/bul0000017
- Wathelet, M., Duhem, S., Vaiva, G., Baubet, T., Habran, E., Veerapa, E., Debien, C., Molenda, S., Horn, M., Grandgenèvre, P., Notredame, C.-E., & D'Hondt, F. (2020). Factors Associated With Mental Health Disorders Among University Students in France Confined During the COVID-19 Pandemic. *JAMA Network Open*, *3*(10), e2025591. doi: 10.1001/jamanetworkopen.2020.25591
- Zugliani, M., Martin-Santos, R., Nardi, A., & Freire, R. (2017). Personality Traits in Panic Disorder Patients With and Without Comorbidities. *The Journal of Nervous and Mental Disease*, *205*(11), 855-858. doi: 10.1097/NMD.0000000000000745