

Comparing the Global Assessment of Functioning (GAF) and the World Health Organization Disability Assessment Schedule (WHODAS) 2.0 in children and adolescents

Francisco R. de la Peña,^{1,✉} Mariana Paulina Escalona,¹ Rosa Elena Ulloa,^{2,✉} Lino Palacios-Cruz,^{1,✉} Juan David Palacio,^{3,✉} Pablo Mayer,^{4,✉} Ricardo Diaz,² Marcos Francisco Rosetti^{1,5,✉}

¹ Instituto Nacional de Psiquiatría Ramón de la Fuente Muñiz, Ciudad de México, México.

² Hospital Psiquiátrico Infantil "Juan N. Navarro", Ciudad de México, México.

³ Facultad de Medicina, Universidad de Antioquia, Medellín Colombia.

⁴ Universidad Autónoma Metropolitana, Toluca, México.

⁵ Instituto de Investigaciones Biomédicas, Universidad Nacional Autónoma de México, Ciudad de México, México.

Correspondence:

Marcos Francisco Rosetti
Unidad Psicopatología y Desarrollo,
Centro de Investigación en Salud
Mental Global, Instituto Nacional
de Psiquiatría Ramón de la Fuente
Muñiz.
Calzada México - Xochimilco 101,
Col. Huipulco, C.P. 14370,
Ciudad de México, México.
Phone: 52+ 55 4161-2485
Email: mrosetti@gmail.com

Received: June 27, 2023

Accepted: November 14, 2023

Citation:

De la Peña, F. R., Escalona, M. P., Ulloa, R. E., Palacios-Cruz, L., Palacio, J. D., Mayer, P., Diaz, R., & Rosetti, M. F. (2024). Comparing the Global Assessment of Functioning (GAF) and the World Health Organization Disability Assessment Schedule (WHODAS) 2.0 in children and adolescents. *Salud Mental*, 47(3), 137-143.

DOI: 10.17711/SM.0185-3325.2024.018



ABSTRACT

Introduction. The DSM-5 replaced the Global Assessment of Functioning (GAF) with the World Health Organization Disability Assessment Questionnaire (WHODAS) 2.0 as a measure of functioning because of the shortcomings of the former. However, further evidence of how GAF and WHODAS 2.0 scores are correlated and how both instruments are associated with sociodemographic and clinical variables, particularly in children and adolescents, is required to support this change. **Objective.** To correlate GAF and WHODAS 2.0 scores in a sample of children and adolescents, and to determine which sociodemographic and clinical variables are associated with the scores of each instrument. **Method.** Using reports obtained from a secondary database analysis of a cross-sectional, multicentric study, we calculated the correlation between WHODAS 2.0 and GAF scores in a clinical sample of children and performed a general linear model analysis to evaluate the association between the sociodemographic and clinical variables with functioning scores. **Results.** Sixty-six participants completed the evaluation. The correlation between WHODAS 2.0 and GAF ($r = -.69$, 95% CI = $[-.82, -.49]$, $p < .001$) was moderate to large and significant. Only poor peer relationships and a higher number of diagnoses were significantly associated with low functioning in both instruments. The results suggest that WHODAS 2.0 and GAF scores reflect different aspects of functioning and disability. **Discussion and conclusion.** Both instruments can provide an accurate assessment of disability/functionality. We propose that, for pediatric cases, WHODAS could provide more information on the self-care domain.

Keywords: Functioning, severity, psychopathology, WHODAS 2.0 interview version, GAF, children and adolescents.

RESUMEN

Introducción. El DSM-5 reemplazó la Evaluación Global del Funcionamiento (GAF) con el Cuestionario de Evaluación de la Discapacidad de la Organización Mundial de la Salud (WHODAS) 2.0 como medida de funcionamiento debido a las deficiencias del primero; sin embargo, mayor evidencia de cómo se correlacionan las puntuaciones GAF y WHODAS 2.0 y cómo ambos instrumentos se asocian con variables sociodemográficas y clínicas, especialmente en niños y adolescentes se necesita para apoyar este cambio. **Objetivo.** Correlacionar las puntuaciones GAF y WHODAS 2.0 en una muestra de niños y adolescentes, y evaluar qué variables sociodemográficas y clínicas están asociadas a las puntuaciones de cada uno de los instrumentos. **Método.** Utilizando informes obtenidos de un análisis de base de datos secundaria de un estudio transversal multicéntrico, calculamos la correlación entre los puntajes WHODAS 2.0 y GAF en una muestra clínica de menores y realizamos un análisis de modelo lineal general para evaluar la asociación entre las variables sociodemográficas y clínicas con puntajes de funcionamiento. **Resultados.** Sesenta y seis participantes completaron la evaluación. La correlación entre WHODAS 2.0 y GAF ($r = -.69$, 95% CI = $[-.82, -.49]$, $p < .001$) fue entre moderada y grande y significativa. Solo las malas relaciones con los compañeros y un mayor número de diagnósticos se asociaron significativamente con un bajo funcionamiento en ambos instrumentos. Estos resultados sugieren que las puntuaciones de WHODAS 2.0 y GAF parecen reflejar diferentes aspectos del funcionamiento y la discapacidad. **Discusión y conclusión.** Ambos instrumentos ofrecen una evaluación certera de la discapacidad/functionality. Para casos pediátricos, el WHODAS 2.0 puede ser más informativo en relación con el auto cuidado.

Palabras clave: Funcionamiento, severidad, psicopatología, WHODAS 2.0 versión de entrevista, GAF, niños y adolescentes.

INTRODUCTION

Some of the changes introduced by the 5th edition of the Diagnostic and Statistical Manual of Mental Disorders (DSM-5) involve the way functioning and disability are assessed. In the DSM-5, the World Health Organization Disability Assessment Schedule (WHODAS) 2.0 replaced the Global Assessment of Functioning (GAF). The DSM-5 Task Force deemed it necessary to replace GAF because of its lack of conceptual clarity and questionable psychometrics in routine practice (American Psychiatric Association, 2013).

WHODAS was created in 1988 as a self-reporting tool to record an individual's perspective on their daily functioning. It provides a standardized score measuring difficulties in functioning in the previous thirty days, regardless of any other diagnosis (Rotter, 2018). Answering WHODAS 2.0 implies a degree of insight that could be challenging for some children. The updated version, WHODAS 2.0, is based on the International Classification of Function (ICF) conceptual framework and captures the level of functioning in six domains of life: cognition, mobility, self-care, relationships, life activities and participation. WHODAS 2.0 places health and disability on a continuum, defining disability as "a decrease in each domain of functioning" (Üstün et al., 2010). In contrast to GAF, the WHODAS 2.0 score does not explicitly consider the severity of mental illness (Gspandl et al., 2018). As WHODAS does not consider the severity of mental disorders as part of the disability/functionality dimension, it allows for the assessment of the latter when there is no correlation with symptom severity (Smith et al., 2011; Von Korff et al., 2011; Söderberg et al., 2005).

In the 3rd edition of the DSM (American Psychiatric Association, 1980), axis V was introduced as a measure of "adaptive functioning" and scored on a 7-point scale ranging from superior to grossly impaired. In the revised 3rd edition of the DSM (American Psychiatric Association, 1987), the GAF scale replaced axis V for the assessment of psychological, social, and occupational functioning. In the 4th edition of the DSM, the GAF scale was extended to a 100-point scale (Pedersen & Karterud, 2012). The GAF scale assesses patient symptoms and overall functional ability on a single hypothetical continuum of mental health-related illness (Gspandl et al., 2018). GAF can be reliable, valid, and sensitive to change over time. However, certain limitations have been widely acknowledged. GAF specifically excludes impairment in functioning due to physical or environmental limitations and scores frequently correlate more with the severity of disorders than with levels of disability (Gold, 2014).

The elimination of GAF and the incorporation of WHODAS 2.0 has posed entailed several challenges due to the differences between these instruments. While the justification for using WHODAS 2.0 to expand the evaluation of global functioning and disability (Reed, 2010) is reason-

able, some questions appear unsuitable for children and adolescents ("Staying by yourself for a few days?" "Sexual activities," "Assuming your household responsibilities" or "Getting your housework done as quickly as needed"). Further research is required to document what exactly is gained and what is lost by this replacement. If they provide different information, the two instruments could be used together in a complementary way.

To date, three studies have been published that evaluate the WHODAS 2.0 and GAF correlation in adult patients. The first evaluated 200 outpatients in Puerto Rico with any diagnosis and reported a positive, significant correlation of .37 between the scores of the two instruments (Martínez-Taboas et al., 2017). The second is a study conducted in the United States evaluating forty-two inpatients and outpatients with schizophrenia, which reported a significant, negative correlation of -.60 using an inverted GAF score. (Gspandl et al., 2018). The third is a Swedish study evaluating 522 patients with any diagnosis, which reported a significant, negative correlation of -.41 (Ramklint et al., 2022). To date, there have been no reports of these correlations in pediatric samples.

Evidence exists that some sociodemographic and clinical variables have been associated with low general functioning and global disability. These two aspects have been linked to lower levels of peer support (Chang et al., 2012), a greater number of comorbid internalizing and externalizing disorders (Dol et al., 2022), perinatal problems, such as preterm birth (Moster et al., 2008), and a history of emotional and sexual abuse (Mullen et al., 1996; Myroniuk et al., 2022). In this respect, it is necessary to evaluate the association between instruments such as the WHODAS 2.0 and GAF and the presence of such variables, particularly in pediatric samples.

We hypothesize that the correlation between WHODAS 2.0 and GAF scores would be low to moderate due to the high variability in severity and functioning evaluation in pediatric samples. Moreover, we consider that high comorbidity, and any kind of abuse would be significantly associated with low functioning in both instruments. The aim of this study was therefore to evaluate the correlation between the WHODAS 2.0 interview version (each domain as well as the total score) and GAF scores and to study the association between clinical and sociodemographic variables and functioning as assessed by the WHODAS 2.0 interview version and GAF in a psychiatric outpatient sample of children and adolescents in Latin America.

METHOD

Participants

The original sample was a convenience sample comprising outpatient children and adolescents ($n = 74$) referred for

medical-psychiatric evaluation in any of the seven clinical sites (see below), from February to August 2016. The study included the subsample of patients who completed WHODAS 2.0 and GAF ($n = 66$).

Raters

Raters were the principal researchers of each clinical site with fifteen years' experience and resident psychiatrists with at least four years' experience in the assessment of pediatric patients. Each clinical site provided three to six different raters. All the raters were trained during a three-day theoretical-practical course when agreements were made in regard to the interview procedures for the K-SADS-PL, GAF and WHODAS 2.0; in the latter, any questions about sexual activity were omitted.

Study design and setting

All participant sites were clinical settings receiving either outpatients, inpatients, or both. The sites included two psychiatric hospitals, a university psychiatric outpatient service in Mexico City, a general hospital with psychiatric service in Aguascalientes, Mexico, a general hospital with psychiatric service in Medellín, Colombia, and two psychiatric hospitals in Santiago de Chile, Chile, and Montevideo, Uruguay.

Measurements

World Health Organization Disability Assessment Schedule (WHODAS) 2.0.

We used the full 36-item self-administered version of WHODAS 2.0, as included in the DSM-5. Studies have reported that the WHODAS 2.0 has high internal consistency (with a Cronbach's alpha of .86), validity and reliability in epidemiological (Kimber et al., 2015) and clinical samples of young adolescents (Hu et al., 2012), and good concurrent validity with other instruments measuring a similar concept of disability (Hernández-Orduña et al., 2017). WHODAS 2.0 questions relate to the difficulties in functioning experienced by the interviewee during the previous thirty days. WHODAS 2.0 scores range from 36 to 180. Due to the low insight reported in psychotic patients (Hernández-Orduña et al., 2017), and the reading difficulties experienced by some children, as instructed by the DSM-5 clinician administration guide, the rater administered WHODAS 2.0 as an interview, simultaneously questioning the child/adolescent and parent/guardian and establishing a consensus answer.

Global Assessment of Functioning (GAF)

The GAF score (1 to 100) provides a classification of the subject's psychological, social, and occupational functioning, ranging from positive mental health to severe psychopathology. The GAF has high internal consistency (with

a Cronbach's alpha of .74) (Söderberg et al., 2005). GAF values can be a single total score or range-type scores (Aas, 2011).

These instruments have opposite directions in their scores. Whereas a higher GAF score suggests higher functioning, a higher WHODAS 2.0 score indicates lower functioning. In this respect, a negative correlation would indicate that both scores correlate in their measure of functioning.

Clinical and psychosocial variables

We used the Kiddie Schedule for Affective Disorders and Schizophrenia (K-SADS-PL-5), a semi-structured diagnostic interview designed to collect detailed sociodemographic and clinical information provided by the parent/guardian and child/adolescent to establish a diagnosis. Through the introductory interview of this instrument, we obtained the full psychosocial information provided by the K-SADS-PL-5, in other words, the participant's status regarding parental divorce, complications at birth, abnormal development, maternal and parental history of mental illness, any kind of abuse and poor peer relationships. All these variables were coded dichotomously, as either present or absent. In the case of abuse, this was scored as present when any type (emotional, sexual, physical, or neglect) was present. Trained clinicians conducted the interview and used all responses to establish a more accurate clinical estimate for the symptoms in each disorder. The number of diagnoses was established after completing all the necessary appendices for each participant. Inter-rater reliability for all diagnoses included in the K-SADS-PL-5 ranged from moderate to good ($\kappa > .7$; de la Peña et al., 2018a; de la Peña et al., 2018b).

Procedure

After receiving a thorough explanation of the research and after signing participation and assent/consent forms, the parent/guardian and the child/adolescent were interviewed for one to four sessions, lasting thirty to 120 minutes each. All participants were evaluated using the Spanish version of the K-SADS-PL-5. The WHODAS 2.0 and GAF were completed at the end of the interview.

Statistical analysis

Descriptive statistics are expressed as measures of means and standard deviation (SD) or frequency and percentages. A Spearman correlation between WHODAS 2.0 and GAF domains and total scores was calculated given the interval nature of the data. A general linear model analysis was performed to evaluate the association between selected sociodemographic and clinical variables with WHODAS 2.0 and GAF functioning scores. Analysis was performed using R (R Core Team, 2021). The level of significance was set at $p < .05$.

Table 1
Psychosocial characteristics and diagnostic profile of the current pediatric sample (n = 66). Psychosocial characteristics were scored dichotomously, and values reflect frequency and percentage of positive answers

Variable	Value	n	(%)
Psychosocial characteristic	Monoparental family	32	(48.4)
	Divorced parents	23	(34.8)
	Developmental delays	7	(10.6)
	Birth delivery complications	30	(45.4)
	Abuse	12	(18.1)
	Bad peer relationships	17	(25.7)
	Mother psychiatric history	15	(22.7)
	Father psychiatric history	16	(24.2)
	Attending School	61	(92.4)
Diagnostic profile	Attention deficit/hyperactivity disorder	25	(37.9)
	Oppositional defiant disorder	18	(27.3)
	Simple phobias	17	(25.6)
	Social anxiety	17	(25.6)
	Intermittent explosive disorder	15	(22.7)
	Major depressive disorder	15	(22.7)
	Substance abuse	2	(3.03)

Ethical considerations

Informed assent and consent were obtained from all child/adolescent and parent/guardian participants in written form. Forms were signed after assenting/consenting to participate, with two witnesses also signing the same form. The authors declare that all procedures contributing to this study comply with the ethical standards of the relevant national and institutional committees on human experimentation and with the Helsinki Declaration of 1975, as revised in 2008. All procedures involving human subjects/patients were approved by the Institutional Review Board of the INPRFM with authorization No. CEI/029/2020.

RESULTS

Sociodemographic and clinical characteristics are shown in Table 1. Most of the total sample ($n = 66$, 62.1% males, 11.4 years, $SD = 3.2$) were children actively attending school (92.4%). The mean number of diagnoses was 3.38 ($SD = 2.78$). More sample details can be reviewed in previous publications (de la Peña et al., 2018a; de la Peña et al., 2018b).

The mean WHODAS 2.0 and GAF scores were 60.48 ($SD = 18.33$) and 59.86 ($SD = 17.10$) respectively. WHODAS mean subdomains yielded the following scores: cognition = 11.38 ($SD = 4.73$), mobility = 6.41 ($SD = 2.08$), self-care = 5.12 ($SD = 1.72$), relationships = 7.5 ($SD = 3.01$), life activities = 16.86 ($SD = 7.68$) and participation = 13.23 ($SD = 5.36$). The correlations between WHODAS 2.0 domains and total scores with GAF were all negative and significant (Table 2).

Regression analysis showed that although several variables contributed significantly to the variance of WHODAS 2.0 ($R^2 = .41$) and GAF ($R^2 = .53$), only the total number of diagnoses and poor peer relationship significantly contributed to the variance of both instruments (Table 3).

DISCUSSION AND CONCLUSION

We found negative and significant correlations between WHODAS 2.0 domains and total scores and GAF. Moreover, we found that only poor peer relationships and a higher number of diagnoses were associated with lower functioning scores on both instruments.

WHODAS 2.0 vs GAF

Some advantages and disadvantages have been described for both instruments. On the one hand, WHODAS 2.0

Table 2
Correlations of World Health Organization Disability Assessment Schedule 2.0 (WHODAS) 2.0 domains and total scores with Global Assessment of Functioning (GAF) scores

Comparisons	r	p	95% CI
WHODAS total score vs. GAF	-.69	< .001	-.81, -.50
WHODAS: cognition vs. GAF	-.50	< .001	-.67, -.29
WHODAS: mobility vs. GAF	-.43	.005	-.61, -.21
WHODAS: self-care vs. GAF	-.27	.034	-.48, -.04
WHODAS: relationships vs. GAF	-.48	.002	-.67, -.27
WHODAS: life activities vs. GAF	-.60	< .001	-.75, -.42
WHODAS: participation vs. GAF	-.55	< .001	-.71, -.35

Table 3
Associations between sociodemographic and clinical variables with World Health Organization Disability Assessment Schedule 2.0 (WHODAS) 2.0 and Global Assessment of Functioning (GAF) scores

Variables	WHODAS 2.0		GAF	
	Est. (S.E.)	<i>p</i>	Est. (S.E.)	<i>p</i>
Age	.98 (.56)	.09	-1.08 (.5)	.03
Gender: Female	-6.45 (3.92)	.11	4.92 (3.46)	.16
Parental divorce+	7.55 (4)	.06	4.22 (3.54)	.24
Birth delivery complications+	.71 (3.68)	.85	-6.55 (3.25)	.05
Abnormal development+	10.56 (5.96)	.08	-6.13 (5.26)	.25
Maternal history of mental illness+	-7.87 (4.96)	.12	6.21 (4.38)	.16
Paternal history of mental illness+	-1.42 (4.29)	.74	-1.02 (3.79)	.79
Abuse+	14.76 (5.13)	.01	-7.73 (4.54)	.09
Bad peer relationships+	13.75 (5.1)	.01	-9.73 (4.5)	.03
Number of diagnoses	1.81 (.69)	.01	-3.12 (.61)	< .001

Note: + Indicates presence vs absence of this characteristic. Significant values are in italics. S.E.= Standard error.

is completed during a semi-structured interview for any disease assessment of patient functioning in which impairment and disability are evaluated over six domains described during the past thirty days. It is reliable, responsive to change, and promoted by the WHO (Üstün et al., 2010; World Health Organization, 2010). Disadvantages include its limited familiarity due to its recent incorporation in the DSM-5, long application times (at least in the 36-item version) and the fact that it mainly uses activities that are meaningful for adults. On the other hand, GAF is completed by clinicians, used in a variety of settings, and has acceptable reliability when raters are experienced and trained (Gspandl et al., 2018). The disadvantages of GAF include the fact that it uses a single measure as an operationalization of more than one clinical phenomenon (psychological symptoms and social and occupational functioning). In addition, its validity diminishes when there is a discrepancy between symptom severity and functioning (Aas, 2011; Pedersen & Karterud, 2012). Finally, WHO-DAS subscales provide a detailed view of functionality, whereas GAF does not identify the specific domain that most affects the patient.

Previous studies of adult psychiatric samples have reported moderate to large significant correlations (Gspandl et al., 2018; Martínez-Taboas et al., 2017; Ramklint et al., 2022) between WHODAS 2.0 and GAF scores. In the current study, the moderate correlations could be explained by the fact that WHODAS 2.0 contains non-specific items, which most children or adolescents with psychopathology can perform without difficulty, such as those in the mobility and self-care domains. In our pediatric sample, these items were rated as “without problems,” leading to a better functioning score, whereas in the GAF evaluation, the clinician integrates functioning with psychiatric disorders,

even though this is a controversial strategy (Smith et al., 2011). In view of this, we would recommend WHODAS 2.0 to evaluate psychiatric patients with a general medical condition that may include mobility, relationships, self-care, or participation disturbances that may not be present in all psychiatric patients. Interestingly, five out of the six domains in WHODAS 2.0 showed a moderate, significant correlation with GAF scores.

Although a WHODAS 2.0 version is available for children and adolescents, it is remarkably similar to the adult version (Scorza et al., 2013) but has not yet been incorporated into the DSM-5 or the DSM-5 revised version. The pediatric version requires further research before it can be widely used.

Functioning vs. sociodemographic and clinical variables

The results of the regression analyses confirmed previous reports regarding the association between sociodemographic and clinical variables and general functioning. In particular, poor peer relationships and a higher number of comorbid disorders were associated with low functioning in the scores of both instruments. In poor peer relationships, both instruments contain items that exclusively assess issues in this domain, which could explain the over-representation. Despite this, social relations in children and adolescents are one of the main pillars of the evaluation of functioning as was seen in the current sample. Children and adolescents spend a great deal of time with their peers who influence their behavior and values. Peer relationships serve as a bridge when adolescents move away from their parents towards independent adult functioning (Rohrbeck, 2003).

As for the number of diagnoses, a growing body of evidence about the commonality of triggers or factors as-

sociated with global psychopathology has led to a more dimensional approach to understanding mental illness. All psychopathological disorders are correlated not only at the disorder level, but also at the spectra level (Wright et al., 2013). Most common psychiatric disorders are therefore unified by a single psychopathology dimension, the p-factor (Caspi et al., 2014). Higher p-factor scores are associated with greater impairment in life and family dysfunction (Caspi & Moffitt, 2018). Previous studies have established that a higher number of diagnoses correlates with poor functioning (Caspi et al., 2014). If a higher number of diagnoses is associated with lower functioning, the dimensional diagnosis approach may need to be more frequently implemented. Variables such as complications at birth and having a history of abuse should be considered in future research to expand and achieve a better understanding of their impact on functioning outcomes.

Limitations

None of the participants in the sample were inpatients with severe disorders, which limits comparisons with other reports. The fact that this was a cross-sectional study meant that no association over time could be reported. The small sample size may have created a statistical bias. The way WHODAS 2.0 was applied limits its comparison with other self-report studies. Since global clinical severity was not evaluated, we were unable to use it as a covariate in the regression analysis.

Conclusion

The results of the current study provide evidence that WHODAS 2.0 and GAF scores appear to reflect similar aspects of functioning, as evidenced by the moderate correlation values. Poor peer relationships and a higher number of diagnoses are associated with lower functioning evaluations with WHODAS 2.0 and GAF. We therefore propose that in the case of younger patients, GAF could be used to complement WHODAS to obtain a full profile of the patient's functioning, particularly in the self-care domain.

Funding

The authors received financial support for the data collection for this research from the Consejo Nacional de Ciencia y Tecnología (Grant No. 23382).

Conflict of interest

The authors declare they have no conflicts of interest.

Acknowledgments

The authors would like to thank Silvia Ortiz, Aurora Jaimes, Marcela Larraguibel, and Laura Viola for their participation in the data collection and all the participants who made this study possible. All the authors declare they have no conflicts of interest.

REFERENCES

- Aas, I. H., (2011). Guidelines for rating global assessment of functioning (GAF). *Annals of General Psychiatry, 10*(1), 1-11. doi: 10.1186/1744-859X-10-2
- American Psychiatric Association. (1980). *Diagnostic and statistical manual: mental disorders* (3rd Ed). American Psychiatric Association.
- American Psychiatric Association. (1987). *Diagnostic and statistical manual: mental disorders: DSM-III-R* (3rd Ed, revised). American Psychiatric Association.
- American Psychiatric Association. (2013). *Diagnostic and statistical manual: mental disorders: DSM-5™* (5th Ed). American Psychiatric Association.
- Caspi, A., & Moffitt, T. E. (2018). All for one and one for all: Mental disorders in one dimension. *American Journal of Psychiatry, 175*(9), 831-844. doi: 10.1176/appi.ajp.2018.17121383
- Caspi, A., Houts, R. M., Belsky, D. W., Goldman-Mellor, S. J., Harrington, H., Israel, S., Meier, M. H., Ramrakha, S., Shalev, I., Poulton, R., & Moffitt, T. E. (2014). The p factor: one general psychopathology factor in the structure of psychiatric disorders? *Clinical Psychological Science, 2*(2), 119-137. doi: 10.1177/2167702613497473
- Chang, E., Eddins-Folensbee, F., & Coverdale, J. (2012). Survey of the prevalence of burnout, stress, depression, and the use of supports by medical students at one school. *Academic Psychiatry, 36*(3), 177-182. doi: 10.1176/appi.ap.11040079
- de la Peña, F. R., Rosetti, M. F., Rodríguez-Delgado, A., Villavicencio, L. R., Palacio, J. D., Montiel, C., Mayer, P. A., Félix, F. J., Larraguibel, M., Viola, L., Ortiz, S., Fernández, S., Jaimes, A., FERIA, M., Sosa, L., Palacios-Cruz, L., & Ulloa, R. E. (2018a). Construct validity and parent-child agreement of the six new or modified disorders included in the Spanish version of the Kiddie Schedule for Affective Disorders and Schizophrenia present and Lifetime Version DSM-5 (K-SADS-PL-5). *Journal of Psychiatric Research, 101*, 28-33.
- de la Peña, F. R., Villavicencio, L. R., Palacio, J. D., Félix, F. J., Larraguibel, M., Viola, L., Ortiz, S., Rosetti, M., Abadi, A., Montiel, C., Mayer, P. A., Fernández, S., Jaimes, A., FERIA, M., Sosa, L., Rodríguez, A., Zavaleta, P., Uribe, D., Galicia, F., ... Ulloa, R. E. (2018b). Validity and reliability of the kiddie schedule for affective disorders and schizophrenia present and lifetime version DSM-5 (K-SADS-PL-5) Spanish version. *BMC Psychiatry, 18*(1), 1-7. doi: 10.1186/s12888-018-1773-0
- Dol, M., Reed, M., & Ferro, M. A. (2022). Internalizing-Externalizing Comorbidity and Impaired Functioning in Children. *Children, 9*(10), 1547. doi: 10.3390/children9101547
- Gold, L. H. (2014). DSM-5 and the assessment of functioning: the World Health Organization Disability Assessment Schedule 2.0 (WHODAS 2.0). *Journal of the American Academy of Psychiatry and the Law Online, 42*(2), 173-181.
- Gspandl, S., Peirson, R. P., Nahhas, R. W., Skale, T. G., & Lehrner, D. S. (2018). Comparing global assessment of functioning (GAF) and World Health Organization disability assessment schedule (WHODAS) 2.0 in schizophrenia. *Psychiatry Research, 259*, 251-253. doi: 10.1016/j.psychres.2017.10.033
- Hernández-Orduña, O., Robles-García, R., Martínez-López, N., Muñoz-Toledo, C., González-Salas, A., Cabello, M., Domínguez-Martínez, T., & Medina-Mora, M. E. (2017). WHODAS and the evaluation of disability among people with mental disorders with and without psychotic symptoms. *Salud Mental, 40*(5), 209-217. doi: 10.17711/SM.0185-3325.2017.027
- Hu, L., Zang, Y.-L., & Li, N. (2012). The applicability of WHODAS 2.0 in adolescents in China. *Journal of Clinical Nursing, 21*(17-18), 2438-2451. doi: 10.1111/j.1365-2702.2012.04126.x
- Kimber, M., Rehm, J., & Ferro, M. A. (2015). Measurement invariance of the WHODAS 2.0 in a population-based sample of youth. *PLoS One, 10*(11), e0142385. doi: 10.1371/journal.pone.0142385
- Martínez-Taboas, A., Medina-Sustache, E., González-Díaz, D. Y., Prats Aparicio, A. C., Garrahan Viejo, A. M., García, J. D. J., Landrau Sálamo, A., & Rodríguez-Cay, J. R. (2017). El WHODAS 2.0 en Puerto Rico: psicometría y su relación con la escala de evaluación de actividad global con pacientes psiquiátricos. *Salud y Sociedad, 8*(1), 82-93. doi: 10.22199/S07187475.2017.0001.00006
- Moster, D., Lie, R. T., & Markestad, T. (2008). Long-term medical and social consequences of preterm birth. *New England Journal of Medicine, 359*(3), 262-273. doi: 10.1056/NEJMoa0706475
- Mullen, P. E., Martin, J. L., Anderson, J. C., Romans, S. E., & Herbison, G. P. (1996). The long-term impact of the physical, emotional, and sexual abuse of children:

- A community study. *Child Abuse & Neglect*, 20(1), 7-21. doi: 10.1016/0145-2134(95)00112-3
- Myroniuk, S., Reitsema, A. M., de Jonge, P., & Jeronimus, B. F. (2022). Specific types of childhood abuse and neglect and profiles of adult emotion dynamics. *PsyArXiv*.
- Pedersen, G., & Karterud, S. (2012). The symptom and function dimensions of the Global Assessment of Functioning (GAF) scale. *Comprehensive Psychiatry*, 53(3), 292-298. doi: 10.1016/j.comppsy.2011.04.007
- R Core Team. (2021). *R: A language and environment for statistical computing*. R Foundation for Statistical Computing, Vienna, Austria. <https://www.R-project.org/>
- Ramklint, M., Söderberg, P., Tungström, S., Nordenskjöld, A., & Hermansson, L. (2022). Validity of the self-rated 36-item World Health Organization Disability Assessment Schedule (WHODAS) 2.0 as a measure of functioning in Swedish psychiatric outpatients. *Nordic Journal of Psychiatry*, 77(3), 276-281. doi: 10.1080/08039488.2022.2097738
- Reed, G. M. (2010). Toward ICD-11: Improving the clinical utility of WHO's International Classification of mental disorders. *Professional Psychology: Research and Practice*, 41(6), 457-464. doi: 10.1037/a0021701
- Rohrbeck, C. A. (2003). Chapter - Peer relationships, adolescence. In *Encyclopedia of primary prevention and health promotion* (pp. 808-812). Boston, MA: Springer.
- Rotter, K. (2018). Valoración de la salud y la discapacidad: WHODAS 2.0. *Rehabilitación Integral*, 13(1), 6-7.
- Scorza, P., Stevenson, A., Canino, G., Mushashi, C., Kanyanganzi, F., Munyanah, M., & Betancourt, T. (2013). Validation of the "World Health Organization disability assessment schedule for children, WHODAS-child" in Rwanda. *PLoS One*, 8(3), e57725. doi: 10.1371/journal.pone.0057725
- Smith, G. N., Ehmann, T. S., Flynn, S. W., MacEwan, G. W., Tee, K., Kopala, L. C., Thornton, A. E., Schenk, C. H., & Honer, W. G. (2011). The assessment of symptom severity and functional impairment with DSM-IV Axis V. *Psychiatric Services*, 62(4), 411-417. doi: 10.1176/appi.ps.62.4.411
- Söderberg, P., Tungström, S., & Armelius, B. Å. (2005). Special section on the GAF: Reliability of Global Assessment of Functioning ratings made by clinical psychiatric staff. *Psychiatric Services*, 56(4), 434-438. doi: 10.1176/appi.ps.56.4.434
- Üstün, T. B., Chatterji, S., Kostanjsek, N., Rehm, J., Kennedy, C., Epping-Jordan, J., Saxena, S., von Korf, M., & Pull, C. (2010). Developing the World Health Organization disability assessment schedule 2.0. *Bulletin of the World Health Organization*, 88(11), 815-823. doi: 10.2471/BLT.09.067231
- Von Korf, M., Katon, W. J., Lin, E. H., Ciechanowski, P., Peterson, D., Ludman, E. J., Young, B., & Rutter, C. M. (2011). Functional outcomes of multi-condition collaborative care and successful ageing: results of randomised trial. *BMJ*, 343, d6612. doi: 10.1136/bmj.d6612
- World Health Organization. (2010). *A conceptual framework for action on the social determinants of health*. Available from: <https://apps.who.int/iris/handle/10665/44489>
- Wright, A. G., Krueger, R. F., Hobbs, M. J., Markon, K. E., Eaton, N. R., & Slade, T. (2013). The structure of psychopathology: toward an expanded quantitative empirical model. *Journal of Abnormal Psychology*, 122(1), 281-294. doi: 10.1037/a0030133